

University of Rajshahi

Rajshahi-6205

Bangladesh.

RUCL Institutional Repository

<http://rulrepository.ru.ac.bd>

Department of Mathematics

MPhil Thesis

2004

Some Topics in Fuzzy Topological Spaces


Hossain, Md. Sahadat

University of Rajshahi

<http://rulrepository.ru.ac.bd/handle/123456789/859>

Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

Some Topics in Fuzzy Topological Spaces



A

*thesis submitted to the Department of Mathematics
University of Rajshahi, Rajshahi – 6205; Bangladesh
for the degree of Master of Philosophy*

in

Mathematics

By

MD. SAHADAT HOSSAIN

Under the supervision of
Professor, Dr. Dewan Muslim Ali
Department of Mathematics
University of Rajshahi
Rajshahi – 6205
Bangladesh



**Dedicated
To My Parents**

D-2436

DOC

514.3

HOS

ACKNOWLEDGEMENTS

Firstly, I give my limitless thanks to the Almighty Allah for giving me strength , endurance and ability to complete this course of study.

It gives me a great pleasure to offer my profound gratitude to my reverend supervisor Dr. Dewan Muslim Ali , Professor , Department of Mathematics , University of Rajshahi , Bangladesh , who taught the topic of the present thesis and provided an excellent guidance with this continuous devotion , endless inspiration and constant encouragement . His invaluable advice, constructive suggestion and willingness to discuss the topic at many odd hours inspite of his busy routine have gone a long way in the successful completion of this thesis in time.

I also wish to express my sincere admiration , appreciation and gratitude to all the teachers of the Department of Mathematics , University of Rajshahi for providing me with invaluable suggestions during the period of my research work.

I want to extend my thanks to the concerned authority of the University of Rajshahi for providing me with a fellowship for the purpose of completing this thesis and thanks are also due to the officers and other staffs of mathematics department for their excellent co-operations.

My sense of fathomless gratitude is to my parents and other members of the family for their unmeasured sacrifices , continuous inspiration and support , that have led me to all the success .

In fine, I am alone responsible for the errors and shortcomings in this study if there be any, I am sorry for that.

Department of Mathematics

(Md . Sahadat Hossain)

December - 2004



No

Dated

CERTIFICATE

I have the pleasure in certifying that the M. Phill. thesis entitled "Some Topics in Fuzzy Topological Spaces" submitted by Md. Sahadat Hossain in fulfillment of the requirement for the degree of M. Phill. in Mathematics, University of Rajshahi, Rajshahi, Bangladesh has been completed under my supervision. I believe that the research work is an original one and it has not been submitted elsewhere for any degree.

I wish him a bright future and every success in life.

Supervisor

A handwritten signature in black ink, appearing to read 'Dewan'.

(Dr. Dewan Muslim Ali)

29.12.04

Professor ,

Department of Mathematics
University of Rajshahi
Rajshahi , Bangladesh

STATEMENT OF ORIGINALITY

I declare that the contents in my M. Phill. thesis entitled " Some Topics in Fuzzy Topological spaces " is original and accurate to the best of my knowledge. I also declare that the materials contained in my research work have not been previously published or written by any person for any degree or diploma.

Rajshahi University

December- 2004

M. Phill. Research Fellow

Md. Sahadat Hossain

23-12-2004

(Md. Sahadat Hossain)

CONTENTS

	Page no.
Introduction	1 – 2
Chapter 1 : PREREQUISITES	3 – 12
1.1. Introduction	3
1.2. Fuzzy set	4
1.3. Mappings and fuzzy subsets introduced mappings	6
1.4. Continuous map, open map and closed map	9
1.5. Subspace topology, base and subbase	10
1.6. Product topology	10
1.7. Good extension	12
CHAPTER 2 : R_0 and R_1 Fuzzy Topological Spaces	13-27
2.1. Introduction	13
2.1.1. R_0 - Fuzzy Topological space	13-19
2.1.1. R_1 - Fuzzy Topological Space	19 – 27
CHAPTER 3: T_0 and T_1 Fuzzy Topological Spaces	28 – 46
3.1. Introduction	28
3.1.1. T_0 – Fuzzy Topological Space	28 – 36
3.2.1. T_1 – Fuzzy Topological Space	36 – 46
CHAPTER 4: T_2 Fuzzy topological Space	47 – 56
4. Introduction	47

	Page no.
4.1. T_2 – Fuzzy Topological Space	47 – 56
CHAPTER 5: Regular Fuzzy Topological Space	57- 66
5. Introduction	57
5.1. Regular- Fuzzy Topological Space	57 – 66
CHAPTER 6: Normal Fuzzy Topological Space	67- 72
6. Introduction	67
6.1. Normal- Fuzzy Topological Space	67- 72
REFERENCES	72 - 80

Introduction

The concept of fuzzy set was introduced by the American Mathematician L . A . Zadeh for the first time in 1965 . This provides a natural framework for generalizing many algebraic and topological concepts in various directions. Accordingly fuzzy groups , fuzzy ideals , fuzzy rings , fuzzy vector space , fuzzy measure , fuzzy topology , fuzzy topological groups, fuzzy topological vector space and many other branches have been developed all over the world during the last four decades . It is still developing in many directions. While reviewing the literature in fuzzy topology , we have seen the gap in separation axiom of fuzzy topological space , the counter part of which in ordinary topological space drew attention of several eminent mathematicians like Azad , K. K. ; Chang , C. L. ; Ali , D. M. ; Wuyts , P. ; Srivastava , A. K. and Lowen , R. etc. We aim to develop of theories of Fuzzy T_0 , Fuzzy T_1 , Fuzzy T_2 , Fuzzy R_0 , Fuzzy R_1 , Fuzzy Regular and Fuzzy Normal spaces analogous to its counter part in ordinary topology . The material of the thesis has been divided into six chapters , a brief scenario of which we present as follows.

Chapter one incorporates some of the basic definitions and results of fuzzy set, fuzzy topology and mappings. These results are ready references for the work in subsequent chapters . Results are stated without proof and can be seen in the papers referred to .

Our work starts from the second chapter. In the second chapter , we have introduced and studied some R_0 and R_1 properties in fuzzy topological spaces and we have established relations among them . Also we have studied some other properties of these concepts.

In the third chapter , We have introduced and studied some T_0 and T_1 properties in fuzzy topological spaces and we have established relations among them . Also we have studied some other properties of these concepts.

In the fourth chapter, we have introduced and studied some T_2 properties in fuzzy topological spaces and we have established relations among them . Also we have studied some other properties of these concepts.

In the fifth chapter, we have introduced and studied some Regular properties in fuzzy topological spaces and we have established relations among them. Also we have studied some other properties of these concepts.

In the sixth chapter, we have introduced and studied some Normal properties in fuzzy topological spaces and we have established relations among them. Also we have studied some other properties of these concepts.

CHAPTER:-1

Prerequisites

1.1. Introduction: - This chapter incorporates concepts and results of the Fuzzy sets, Fuzzy topological spaces and Fuzzy product topological spaces which are to be used as ready references for understanding the subsequent chapters. Most of the results are quoted from various research papers. Through the sequel, we make use the following notations.

Λ : Index set.

$I = [0, 1]$: Closed unit interval.

$I_1 = [0, 1)$: Right open unit interval.

$I_0 = (0, 1]$: Left open unit interval.

u, v, w, \dots : Fuzzy sets.

(X, t) : Fuzzy topological space.

(X, T) : General topological space.

$\prod_{i \in \Lambda} X_i$: Usual product of X_i .

$(X, t_1 \times t_2)$: Product of fuzzy topologies t_1 and t_2 on the set X .

$I_\alpha(t) = \{ u^{-1}(\alpha, t) : u \in t \}, \alpha \in I_1$ General topology on X .

1.2. This thesis deals with the study of fuzzy topological spaces. To present our work in a systematic way in this thesis, we consider in this chapter, various concepts and results on fuzzy sets and fuzzy topological spaces scattered in various research papers. For this we start with .

1.2.1. Definition:- Let X be a nonempty set and A be a subset of X . The function

$1_A : X \longrightarrow [0, 1]$ defined by

$$\begin{aligned} 1_A(x) &= 1 && \text{if } x \in A \\ &= 0 && \text{if } x \notin A \end{aligned}$$

is called the characteristic function of A .

1.2.2. Definition [94]:- A function u from X into the unit interval I is called a fuzzy set in X .

For every $x \in X$, $u(x) \in I$ is called the grade of membership (g.m.f) of x in u . Some

authors say that u is a fuzzy subset of X instead of saying that u is a fuzzy set in X .

1.2.3. Definition [56]:- A fuzzy subset is empty if and only if grade of membership is identically zero in X . It is denoted by 0 .

1.2.4. Definition [56] :- A fuzzy subset is whole if and only if its grade of membership is identically one in X . It is denoted by 1 .

1.2.5. Definition [94] :- Let X be a set and u and v be two fuzzy subsets of X . Then u is said to be subset of v if $u(x) \leq v(x)$ for every $x \in X$. It is denoted by $u \subseteq v$.

1.2.6. Definition [94] :- Let X be a set and u and v be two fuzzy subsets of X . Then u is said to be equal to v if and only if $u(x) = v(x)$ for every $x \in X$. It is denoted by $u = v$.

1.2.7. Definition [94] :- Let X be a set and u and v be two fuzzy subsets of X . Then u is said to be the complement of v if $v(x) = 1 - u(x)$, for every $x \in X$. It is denoted by u^c . Obviously $(u^c)^c = u$.

1.2.8. Definition [15] :- Let X be a set and u and v be two fuzzy subsets of X . Then the union w of u and v is a fuzzy subset of X , written as $w = u \cup v$ which is defined by

$$w(x) = (u \cup v)(x) = \max \{ u(x), v(x) \} \text{ for every } x \in X.$$

In general, if Λ be an index set and $A = \{ u_i : i \in \Lambda \}$ be a family of fuzzy sets of X then the union $\cup u_i$ is defined by

$$(\cup u_i)(x) = \sup \{ u_i(x) : i \in \Lambda \}, x \in X.$$

1.2.9. Definition [15] :- Let X be a set and u and v be two fuzzy subsets of X . Then the intersection m of u and v is a fuzzy subset of X , written as $m(x) = (u \cap v)(x) = \min \{u(x), v(x)\}, \forall x \in X$, and $(\cap u_i)(x) = \inf \{u_i(x) : i \in \Lambda\}, x \in X$, where $\{ u_i, i \in \Lambda \}$.

1.2.10. Definition :- Let X be a set u and v be two fuzzy subsets of X . Then the difference of u and v is defined by $u - v = u \cap v^c$.

Laws of the algebra of fuzzy sets :

As in ordinary set theory, idempotent laws, associative law, commutative law, distributive laws, identity law, demorgan's laws hold in the case of fuzzy sets also. But the complement laws are not necessarily true. For example, if $X = \{ a, b, c \}$ and u is a fuzzy subset of X where is defined by

$$u = \{ (a, .2), (b, .7), (c, 1) \},$$

$$\text{then } u^c = \{ (a, .8), (b, .3), (c, 0) \}$$

$$\text{so } u \cup u^c = \{ (a, .8), (b, .7), (c, 1) \} \neq 1,$$

$$u \cap u^c = \{ (a, .2), (b, .3), (c, 0) \} \neq 0.$$

Also in ordinary set theory $U \cap V = \phi$ iff $U \subset V^c$. But in fuzzy subsets reverse is not necessary true. For example if

$$v = \{ (a, .6), (b, .2), (c, 0) \} \text{ then } u \subset v^c,$$

$$u \cap v = \{ (a, .2), (b, .2), (c, 0) \} \neq 0.$$

1.3. Mapping and fuzzy subsets induced by mappings,

1.3.1. Definition [15] :- Let f be a mapping from a set X into a set Y and u be a fuzzy subset of X . Then f and u induced a fuzzy subset v of Y defined by

$$v(y) = \sup \{ u(x) \} \text{ if } f^{-1}[\{y\}] \neq \phi, x \in X \\ = 0 \text{ otherwise .}$$

1.3.2. Definition [15] :- Let f be a mapping from a set X into Y and v be a fuzzy subset of Y . Then the inverse of v written as $f^{-1}(v)$ is a fuzzy subset of X and is defined by

$$f^{-1}(v)(x) = v(f(x)), \text{ for } x \in X.$$

We now mention some properties of fuzzy subsets induced by mappings .

Let f be a mapping from X into Y , u be a fuzzy subset of X and v be a fuzzy subset of Y .

Then the following properties are true [15].

(a) $f^{-1}(v^c) = (f^{-1}(v))^c$ for any fuzzy subset v of Y .

(b) $f(u^c) = (f(u))^c$ for any fuzzy subset u of X .

(c) $v_1 \subset v_2 \Rightarrow f^{-1}(v_1) \subset f^{-1}(v_2)$, where v_1 and v_2 are two fuzzy subsets of Y .

(d) $u_1 \subset u_2 \Rightarrow f(u_1) \subset f(u_2)$, where u_1 and u_2 are two fuzzy subsets of X .

(e) $v \supset f(f^{-1}(v))$, for any fuzzy subset v of Y .

(f) $u \subset f^{-1}(f(u))$, for any fuzzy subset u of X .

(g) Let f be a function from X into Y and g be a function from Y into Z . Then $(g \circ f)^{-1}(w) = f^{-1}(g^{-1}(w))$, for any fuzzy subset w in Z , where $(g \circ f)$ is the composition of g and f .

1.3.3. Definition [56] :- A fuzzy point in X is a special type of fuzzy set in X with membership function $p(x) = r, p(y) = 0, \forall y \neq x$, where $0 < r < 1$. This fuzzy point is said to have support x and value r and this point is denoted by x_r or $r1_x$.

1.3.4. Definition [56] :- A fuzzy point p is said to belong a fuzzy set u in X ($p \in u$) if and only if $p(x) < u(x)$ and $p(y) \leq u(y) \forall y \neq x$. Evidently, every fuzzy set u can be expressed as the union of all the fuzzy points which belong to u .

1.3.5. Definition [56] :- Two fuzzy sets u and v in X are said to be intersected if and only if there exist a point $x \in X$ such that $(u \cap v)(x) \neq 0$. In this case we say that u and v intersect at x .

1.3.6. Definition [89] :- Let $I = [0, 1]$, X be a non empty set, and I^X be the collection of all mappings from X into I , ie the class of all fuzzy sets in X . A fuzzy topology on X is defined as a family t of members of I^X , satisfying the following conditions.

- (i) $1, 0 \in t$,
- (ii) If $u_i \in t$ for each $i \in \Lambda$, then $\cup_{i \in \Lambda} u_i \in t$.
- (iii) If $u_1, u_2 \in t$ then $u_1 \cap u_2 \in t$.

The pair (X, t) is called a fuzzy topological space (fts, in short) and members of t are called t -open (or simply open) fuzzy sets. A fuzzy set v is called a t -closed (or closed) fuzzy set if $1 - v \in t$.

1.3.7. Example :- Let $X = \{a, b, c, d\}, t = \{0, 1, u, v\}$,

where $1 = \{(a, 1), (b, 1), (c, 1), (d, 1)\}$

$0 = \{(a, 0), (b, 0), (c, 0), (d, 0)\}$

$$u = \{ (a, .2), (b, .5), (c, .7), (d, .9) \}$$

$$v = \{ (a, .3), (b, .5), (c, .8), (d, .95) \}$$

Then (X, t) is a fuzzy topological space .

1.3.8. Definition [50] :- (According to lowen) A fuzzy topology on non empty set X is a collection t of fuzzy subsets of X such that

(i) all constant fuzzy subsets of X belong to t .

(ii) t is closed under formation of fuzzy union of arbitrary collection of members of t .

(iii) t is closed under formation of intersection of finite collection of members of t .

1.3.9. Definition [56] :- Let u be a fuzzy set in (X, t) . The interior of u is defined as the union of all t – open sets contained in u . It is denoted by u° . Evidently u° is the largest open fuzzy set contained in u and $(u^\circ)^\circ = u^\circ$.

1.3.10. Definition [56] :- The intersection of all the t – closed set containing u is called the closure of u denoted by \bar{u} . Obviously \bar{u} is the smallest closed set containing u and $\bar{\bar{u}} = \bar{u}$.

1.3.11. Definition [84] :- A fuzzy set n in a t fs (X, t) is called a neighborhood of a point $x \in X$, if and only if there exist $u \in t$ such that $u \subseteq n$ and $u(x) = n(x) > 0$.

1.3.12. Example :- Let us consider the example 1.3.7 ,

and $n = \{ (a, .5), (b, .6), (c, .8), (d, .9) \}$. Now n is a neighborhood of $d \in X$. Since $u \in t$ such that $u \subseteq n$ and $u(d) = n(d) > 0$, n is a neighborhood of d . Similarly

$n_1 = \{ (a, .7), (b, .6), (c, .75), (d, .9) \}$ is a neighborhood of d . we denoted the family of all neighborhoods of x by N_x .

1.3.13. Definition [56] :-A fuzzy set u in a fts (X, t) is called a neighborhood of a fuzzy point x_r if and only if there exist a fuzzy set $u_1 \in t$ such that $x_r \in u_1 \subseteq u$. A neighborhood u is called an open neighborhood if u is open. The family consisting of all the neighborhoods of x_r is called the system of x_r .

1.4. Continuous map, open and closed map.

1.4.1. Definition [57] :- The function $f: (X, t) \longrightarrow (Y, s)$ is called fuzzy continuous if and only if for every $v \in s$, $f^{-1}(v) \in t$, the function f is called fuzzy homeomorphif if and only if f is bijective and both f and f^{-1} are fuzzy continuous.

1.4.2. Definition [53] :- The function $f: (X, t) \longrightarrow (Y, s)$ is called fuzzy open if and only if for each open fuzzy set u in (X, t) , $f(u)$ is open fuzzy set in (Y, s) .

1.4.3. Definition [57] :- The function $f: (X, t) \longrightarrow (Y, s)$ is called fuzzy closed if and only if for each closed fuzzy set u in (X, t) , $f(u)$ is closed fuzzy set in (Y, s) .

1.4.4. Proposition ([57] Theorem 1.1) :- Let $f: (X, t) \longrightarrow (Y, s)$ be a fuzzy continuous function, then the following properties hold :

- (i) For every s – closed v , $f^{-1}(v)$ is t – closed.
- (ii) For each fuzzy point p in X and each neighborhood u of $f(p)$, then there exist a neighborhood v of p such that $f(v) \subseteq u$.
- (iii) For any fuzzy set u in X , $f(\overline{u}) \subseteq \overline{f(u)}$.
- (iv) For any fuzzy set v in Y , $\overline{f^{-1}(v)} \subseteq f^{-1}(\overline{v})$.

1.4.5. Proposition ([53] Theorem 3.1) :- Let $f: (X, t) \longrightarrow (Y, s)$ be a fuzzy open function , then the following properties hold:

(i) $f(u^\circ) \subseteq (f(u))^\circ$, for each fuzzy set u in X .

(ii) $(f^{-1}(v))^\circ \subseteq f^{-1}(v^\circ)$, for each fuzzy set v in Y .

1.4.6. Proposition ([53] Theorem 1.5) :- Let $f: (X, t) \longrightarrow (Y, s)$ be a function. Then f is closed if and only if $\overline{f(u)} \subseteq f(\overline{u})$ for each fuzzy set u in X .

1.5. Subspace topology, Base and Subbase.

1.5.1. Definition [56] :- Let (X, t) be a fuzzy topological space and A be an ordinary subset of X . The class $t_A = \{ u / A : u \in t \}$ determines a fuzzy topology on A . This topology is called the subspace fuzzy topology on A .

1.5.2. Definition [89] :- Let (X, t) be a fuzzy topological space . A subfamily B of t is a base for t if and only if each member of t can be express as the union of some members of B .

1.5.3. Definition[89] :- Let (X, t) be a fuzzy topological space . A subfamily S of t is a sub-base for t if and only if the family of finite intersection of members of S forms a base for t .

1.6. Product topology .

1.6.1. Definition [9] :- If u_1 and u_2 are two fuzzy subsets of X and Y respectively then the Cartesian product $u_1 \times u_2$ of two fuzzy subsets u_1 and u_2 is a fuzzy subsets of $X \times Y$ defined by $(u_1 \times u_2)(x, y) = \min(u_1(x), u_2(y))$, for each pair $(x, y) \in X \times Y$.

1.6.2. Definition [47] :- Let $\{ X_i, i \in \Lambda \}$, be any class of sets and let X denoted the Cartesian product of these sets, ie $X = \prod_{i \in \Lambda} X_i$. Note that X consists of all points $p = \langle a_i, i \in \Lambda \rangle$, where $a_i \in X_i$. Recall that, for each $j_0 \in \Lambda$, we define the projection π_{j_0} from the product set X to the coordinate space X_{j_0} . ie $\pi_{j_0} : X \longrightarrow X_{j_0}$ by $\pi_{j_0} (\langle a_i : i \in \Lambda \rangle) = a_{j_0}$,

These projections are used to defined the product topology.

1.6.3. Definition [89] :- If (X_1, t_1) and (X_2, t_2) be two fuzzy topological space and $X = X_1 \times X_2$ be the usual product and t be the coarsest fuzzy topology on X , then each projection $\pi_i : X \longrightarrow X_i, i = 1, 2$. is fuzzy continuous. The pair (X, t) is called the product space of the fuzzy topological spaces (X_1, t_1) and (X_2, t_2) .

1.6.4. Proposition ([9] Theorem 3.6) :- If u is a fuzzy subset of a fuzzy topological space (X, t_1) and v is a fuzzy subsets of a fuzzy topological space (Y, t_2) , then $\overline{u \times v} \subseteq \overline{u} \times \overline{v}$.

1.6.5. Definition [88] :- Let $\{ X_\alpha \}_{\alpha \in \Lambda}$ be a family of nonempty sets. Let $X = \prod_{\alpha \in \Lambda} X_\alpha$ be the usual product of X_α 's and let π_α be the projection from X into X_α . Further assume that each X_α is an fts with fuzzy topology t_α . Now the fuzzy topology generated by $\{ \pi_\alpha^{-1}(b_\alpha) : b_\alpha \in t_\alpha, \alpha \in \Lambda \}$ as a sub basis, is called the product fuzzy topology on X . Clearly if w is a basis element in the product, then there exist $\alpha_1, \alpha_2, \dots, \alpha_n \in \Lambda$ such that

$$w(x) = \min \{ b_\alpha(x_\alpha) : \alpha = 1, 2, 3, \dots, n \}, \text{ where } x = (x_\alpha)_{\alpha \in \Lambda} \in X.$$

1.7 Good extension :

1.7.1. Definition [71] :- Let f be a real valued function on a topological space . If $\{ x : f(x) > \alpha \}$ is open for every real α , then f is called lower semi continuous function .

1.7.2. Definition [50] :- Let X be a nonempty set and T be a topology on X . Let $t = \omega(T)$ be the set of all lower semi continuous (lsc) functions from (X, T) to I (with usual topology). Thus $\omega(T) = \{ u \in I^X : u^{-1}(\alpha, 1] \in T \}$ for each $\alpha \in I_1$. It can be shown that $\omega(T)$ is a fuzzy topology on X .

Let P be the property of a topological space (X, T) and FP be its fuzzy topological analogue. Then FP is called a 'good extension' of P " iff the statement (X, T) has P iff $(X, \omega(T))$ has FP " holds good for every topological space (X, T) .

Chapter : -2

R_0 and R_1 Fuzzy Topological Spaces

2.1 Introduction :-

In this chapter, we introduce and study some R_0 and R_1 properties in fuzzy topological spaces and obtain their several features .

2.1.1. Definition :-

Let (X, t) be fuzzy topological space and $\alpha \in I_1$.

(a) (X, t) is an $\alpha - R_0$ (i) space $\Leftrightarrow \forall x, y \in X, x \neq y$, whenever $\exists u \in t$ with $u(x) = 1$ and $u(y) \leq \alpha$, then $\exists v \in t$ with $v(x) \leq \alpha$ and $v(y) = 1$.

(b) (X, t) is an $\alpha - R_0$ (ii) space $\Leftrightarrow \forall x, y \in X, x \neq y$, whenever $\exists u \in t$ with $u(x) = 0$ and $u(y) > \alpha$, then $\exists v \in t$ with $v(x) > \alpha$ and $v(y) = 0$.

(c) (X, t) is an $\alpha - R_0$ (iii) space $\Leftrightarrow \forall x, y \in X, x \neq y$, whenever $\exists u \in t$ with $0 \leq u(x) \leq \alpha < u(y) \leq 1$, then $\exists v \in t$ with $0 \leq v(y) \leq \alpha < v(x) \leq 1$.

(d) (X, t) is an R_0 (iv) space $\Leftrightarrow \forall x, y \in X, x \neq y$, whenever $\exists u \in t$ with $u(x) < u(y)$, then $\exists v \in t$ with $v(x) > v(y)$.

2.1.2. Lemma:- Show that the properties $\alpha - R_0$ (i), $\alpha - R_0$ (ii), $\alpha - R_0$ (iii) and R_0 (iv) are all independent .

Proof:- For this , we give some examples .

2.1.3. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1$, $u(y) = 0$ and $v(x) = 0.5$, $v(y) = 1$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$, it is found that (X, t) is $\alpha - R_0(i)$, but (X, t) is not $\alpha - R_0(ii)$.

2.1.4. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0$, $u(y) = 1$ and $v(x) = 0.6$, $v(y) = 0$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.5$, we see that (X, t) is $\alpha - R_0(ii)$, but (X, t) is not $\alpha - R_0(i)$.

2.1.5. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1$, $u(y) = 0$ and $v(x) = 0.2$, $v(y) = 0.7$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.5$, it is easy to see that (X, t) is $\alpha - R_0(iii)$, but (X, t) is not $\alpha - R_0(i)$ and (X, t) is not $\alpha - R_0(ii)$.

2.1.6. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1$, $u(y) = 0$ and $v(x) = 0.2$, $v(y) = 0.4$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.5$, it is clear that (X, t) is $\alpha - R_0(iv)$, but (X, t) is not $\alpha - R_0(i)$, (X, t) is not $\alpha - R_0(ii)$ and (X, t) is not $\alpha - R_0(iii)$.

2.1.7. Example :- Let $X = \{ x, y, z \}$ and $u, v, w \in I^X$, where u, v, w are defined by $u(x) = 1$, $u(y) = 1$, $u(z) = 0$ and $v(x) = 0$, $v(y) = 0$, $v(z) = 1$ and $w(x) = 0.9$, $w(y) = 0.5$, $w(z) = 0$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, w, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$, it can be shown that (X, t) is $\alpha - R_0(i)$ and (X, t) is $\alpha - R_0(ii)$. It is observe that (X, t) is not $\alpha - R_0(iii)$, and (X, t) is not $R_0(iv)$, Since $w(x) > \alpha \geq w(y)$ but there does not exist $q \in t$ such that $q(x) \leq \alpha < q(y)$.

2.1.8. Example :- Let $X = \{ x, y, z \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.8$, $u(y) = 0.4$, $u(z) = 0.3$, and $v(x) = 0.3$, $v(y) = 0.8$, $v(z) = 0.2$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.5$ it can be shown that (X, t) is $\alpha - R_0(\text{iii})$, but (X, t) is not $R_0(\text{iv})$, since $u(y) > u(z)$ but we have no $q \in t$ such that $q(y) < q(z)$.

This completes the proof.

2.1.9. Theorem :- Prove that (X, t) is $0 - R_0(\text{ii}) \Leftrightarrow (X, t)$ is $0 - R_0(\text{iii})$.

Proof:- Let (X, t) be $0 - R_0(\text{ii})$ space. We shall prove that (X, t) is $0 - R_0(\text{iii})$. Let $x, y \in X$ with $x \neq y$ and $u \in t$ such that $0 \leq u(x) \leq 0 < u(y) \leq 1$ ie $u(x) = 0, u(y) > 0$.

Since (X, t) is $0 - R_0(\text{ii})$, then $\exists v \in t$ such that $v(x) > 0, v(y) = 0$. Then it is clear that $0 \leq v(y) \leq 0 < v(x) \leq 1$. Hence (X, t) is $0 - R_0(\text{iii})$.

Conversely, suppose that (X, t) is $0 - R_0(\text{iii})$. We shall prove that (X, t) is $0 - R_0(\text{ii})$. Let $x, y \in X$ with $x \neq y$ and $u \in t$ such that $u(x) = 0$ and $u(y) > 0$. It can be written as $0 \leq u(x) \leq 0 < u(y) \leq 1$. Since (X, t) is $0 - R_0(\text{iii})$, $\exists v \in t$ such that $0 \leq v(y) \leq 0 < v(x) \leq 1$ ie $v(y) = 0$ and $v(x) > 0$. Hence (X, t) is $0 - R_0(\text{ii})$.

This completes the proof.

2.1.10. Theorem:- Let (X, t) be a fuzzy topological space and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$, then

- (a) (X, t) is $\alpha - R_0(\text{iii}) \Leftrightarrow (X, I_\alpha(t))$ is R_0 .
- (b) (X, t) is $\alpha - R_0(\text{i}) \not\Leftrightarrow (X, I_\alpha(t))$ is R_0 .
- (c) (X, t) is $\alpha - R_0(\text{ii}) \not\Leftrightarrow (X, I_\alpha(t))$ is R_0 .
- (d) (X, t) is $R_0(\text{iv}) \not\Leftrightarrow (X, I_\alpha(t))$ is R_0 .

Proof :- Let (X, t) be $\alpha - R_0$ (iii). We shall prove that $(X, I_\alpha(t))$ is R_0 . Let $x, y \in X$, $x \neq y$ and $M \in I_\alpha(t)$ with $x \in M, y \notin M$ or $x \notin M, y \in M$. Suppose $x \in M, y \notin M$. We can write $M = u^{-1}(\alpha, 1]$, where $u \in t$. Then we have $u(x) > \alpha, u(y) \leq \alpha$, ie $0 \leq u(y) \leq \alpha < u(x) \leq 1$. Since (X, t) is $\alpha - R_0$ (iii), for $\alpha \in I_1, \exists v \in t$ such that $0 \leq v(x) \leq \alpha < v(y) \leq 1$, ie $v(x) \leq \alpha, v(y) > \alpha$. It follows that $x \notin v^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and also $v^{-1}(\alpha, 1] \in I_\alpha(t)$. Hence it is clear that $(X, I_\alpha(t))$ is R_0 .

Conversely, suppose that $(X, I_\alpha(t))$ is R_0 . We shall prove that (X, t) is $\alpha - R_0$ (iii). Let $x, y \in X, x \neq y$ and $u \in t$ with $0 \leq u(x) \leq \alpha < u(y) \leq 1$, ie $u(x) \leq \alpha, u(y) > \alpha$, it follows that $x \notin u^{-1}(\alpha, 1], y \in u^{-1}(\alpha, 1]$, and $u^{-1}(\alpha, 1] \in I_\alpha(t)$. Since $(X, I_\alpha(t))$ is R_0 , then $\exists M \in I_\alpha(t)$ such that $x \in M, y \notin M$. We can write $M = v^{-1}(\alpha, 1]$ where $v \in t$, it follows that $v(x) > \alpha, v(y) \leq \alpha$, ie $0 \leq v(y) \leq \alpha < v(x) \leq 1$. Hence it is clear that (X, t) is $\alpha - R_0$ (i), ie (a) is proved.

Now, we give some examples.

2.1.11. Example :- Let $X = \{x, y, z\}$ and $u, v \in I^X$ where u, v are defined by $u(x) = 1, u(y) = 0, u(z) = 0.8$ and $v(x) = 0, v(y) = 1, v(z) = 0.7$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$, it is clear that (X, t) is $\alpha - R_0$ (i). Now $I_\alpha(t) = \{X, \Phi, \{x, z\}, \{y, z\}, \{z\}\}$. It is clear that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is not R_0 space, since $y, z \in X, y \neq z$ and $\{x, z\} \in I_\alpha(t)$, with $z \in \{x, z\}, y \notin \{x, z\}$, but there is no $U \in I_\alpha(t)$ with $z \notin U, y \in U$.

2.1.12. Example :- Let $X = \{x, y, z\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.3, u(y) = 0, u(z) = 0.8$, and $v(x) = 0.8, v(y) = 1, v(z) = 0$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$, it is clear that (X, t) is $\alpha - R_0$ (ii) and (X, t) is also R_0 (iv). Now $I_\alpha(t) = \{X, \Phi, \{z\}, \{y\}, \{y, z\}\}$. It is clear

that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is not R_0 space, since $x, y \in X, x \neq y$ and $\{y\} \in I_\alpha(t)$ with $x \notin \{y\}, y \in \{y\}$, but there is no $U \in I_\alpha(t)$ with $x \in U, y \notin U$.

2.1.13. Example :- Let $X = \{x, y\}$ and $u, v, w \in I^X$, where u, v, w are defined by $u(x) = 1, u(y) = 0, v(x) = 0.4, v(y) = 0.9, w(x) = 0.7, w(y) = 0.3$. Consider the fuzzy topology t on X generated by $\{0, u, v, w, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$, it is clear that (X, t) is not α - R_0 (i) and (X, t) is not α - R_0 (ii). Now $I_\alpha(t) = \{X, \Phi, \{x\}, \{y\}\}$. Then we see that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is R_0 .

2.1.14. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.4, u(y) = 0.5, v(x) = 0.3, v(y) = 0.4$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constant}\}$. For $\alpha = 0.6$, we ~~that~~ ^{see} that (X, t) is not α - R_0 (iv). Now $I_\alpha(t) = \{X, \Phi\}$. Then $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is R_0 .

This completes the proof.

2.1.15. Theorem :-

Let (X, T) be a topological space. Then (X, T) is R_0 , iff $(X, \omega(T))$ is α - R_0 (p), where $p = i, ii, iii, iv$.

Proof :- Let $(X, \omega(T))$ be α - R_0 (i). We shall prove that (X, T) is R_0 . Let $x, y \in X$ with $x \neq y$ and $U \in T$ with $x \in U, y \notin U$. But by the definition of $lsc, 1_U \in \omega(T)$ and $1_U(x) = 1, 1_U(y) = 0$. Now we have $1_U \in \omega(T)$ with $1_U(x) = 1, 1_U(y) \leq \alpha$. Since $(X, \omega(T))$ is α - R_0 (i), for $\alpha \in I_1, \exists v \in \omega(T)$ such that $v(x) \leq \alpha, v(y) = 1$. Then $x \notin v^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ as $v(x) \leq \alpha, v(y) = 1$, and also $\exists v^{-1}(\alpha, 1] \in T$. Hence it is clear that (X, T) is R_0 -space.

Conversely, suppose that (X, T) be a R_0 -space. We shall prove that $(X, \omega(T))$ is $\alpha - R_0$ (i). Let $x, y \in X$ with $x \neq y$ and there exist $u \in \omega(T)$ such that $u(x) = 1$, $u(y) \leq \alpha$. Then $x \in u^{-1}(\alpha, 1]$, $y \notin u^{-1}(\alpha, 1]$ as $u(x) = 1$, $u(y) \leq \alpha$, and it is clear that $u^{-1}(\alpha, 1] \in T$. Since (X, T) is R_0 , then $\exists V \in T$ such that $x \notin V$, $y \in V$, but $1_V \in \omega(T)$ and $1_V(x) = 0$, $1_V(y) = 1$, ie $\exists 1_V \in \omega(T)$ such that $1_V(x) \leq \alpha$, $1_V(y) = 1$. Hence it is clear that $(X, \omega(T))$ is $\alpha - R_0$ (i).

Hence (X, T) is $R_0 \Leftrightarrow (X, \omega(T))$ is $\alpha - R_0$ (i).

In the same way we can prove that

(X, T) is $R_0 \Leftrightarrow (X, \omega(T))$ is $\alpha - R_0$ (ii).

(X, T) is $R_0 \Leftrightarrow (X, \omega(T))$ is $\alpha - R_0$ (iii).

(X, T) is $R_0 \Leftrightarrow (X, \omega(T))$ is R_0 (iv).

Thus it is seen that $\alpha - R_0$ (p) is a good extension of its topological counter part (p = i, ii, iii, iv).

This completes the proof.

2.1.16. Theorem :- Let (X, t) be a fuzzy topological space and $A \subseteq X$,

$t_A = \{ u/A : u \in t \}$, then

(a) (X, t) is an $\alpha - R_0$ (i) $\Rightarrow (A, t_A)$ is an $\alpha - R_0$ (i).

(b) (X, t) is an $\alpha - R_0$ (ii) $\Rightarrow (A, t_A)$ is an $\alpha - R_0$ (ii).

(c) (X, t) is an $\alpha - R_0$ (iii) $\Rightarrow (A, t_A)$ is an $\alpha - R_0$ (iii).

(d) (X, t) is an R_0 (iv) $\Rightarrow (A, t_A)$ is an R_0 (iv).

Proof :- First suppose that (X, t) is $\alpha - R_0$ (i). We shall prove that (A, t_A) is $\alpha - R_0$ (i).

Let $x, y \in A$, with $x \neq y$ and $u \in t_A$ such that $u(x) = 1$, $u(y) \leq \alpha$, then also $x, y \in X$, $x \neq y$. But we can write $u = w/A$ where $w \in t$ and hence $w(x) = 1$, $w(y) \leq \alpha$. Since

(X, t) is $\alpha - R_0(i)$, then $\exists m \in t$ such that $m(x) \leq \alpha$, $m(y) = 1$. But from the definition $m/A \in t_A$, for every $m \in t$ and $m/A(x) \leq \alpha$, $m/A(y) = 1$. Thus (A, t_A) is $\alpha - R_0(i)$.

similarly (b), (c) and (d) can be proved.

This completes the proof.

2.1.17. Theorem :- Given (X_i, t_i) , $i \in \Lambda$ be fuzzy topological spaces and $X = \prod_{i \in \Lambda} X_i$ and t be a product fuzzy topology on X . Then ;

(a) $\forall i \in \Lambda$, (X_i, t_i) is $\alpha - R_0(i) \Leftrightarrow (X, t)$ is $\alpha - R_0(i)$.

(b) $\forall i \in \Lambda$, (X_i, t_i) is $\alpha - R_0(ii) \Leftrightarrow (X, t)$ is $\alpha - R_0(ii)$.

(c) $\forall i \in \Lambda$, (X_i, t_i) is $\alpha - R_0(iii) \Leftrightarrow (X, t)$ is $\alpha - R_0(iii)$.

(d) $\forall i \in \Lambda$, (X_i, t_i) is $R_0(iv) \Leftrightarrow (X, t)$ is $R_0(iv)$.

Proof :- Let (X_i, t_i) , $i \in \Lambda$ be $\alpha - R_0(i)$. We shall prove that (X, t) is $\alpha - R_0(i)$.

Let $x, y \in X$, with $x \neq y$ and $u \in t$ such that $u(x) = 1$, $u(y) \leq \alpha$. But we have $u(x) = \min\{u_i(x_i) : i \in \Lambda\}$ and $u(y) = \min\{u_i(y_i) : i \in \Lambda\}$ and hence we can find an $u_i \in t_i$ and $x_i \neq y_i$ such that $u_i(x_i) = 1$ and $u_i(y_i) \leq \alpha$. Since (X_i, t_i) , $i \in \Lambda$ is $\alpha - R_0(i)$, $\alpha \in I_1$, then $\exists v_i \in t_i$, such that $v_i(x_i) \leq \alpha$, $v_i(y_i) = 1$. But we have $\pi_i(x) = x_i$ and $\pi_i(y) = y_i$ and hence $v_i(\pi_i(x)) \leq \alpha$, $v_i(\pi_i(y)) = 1$. It follows that $\exists u_i \circ \pi_i \in t$ such that $(u_i \circ \pi_i)(x) \leq \alpha$, $(u_i \circ \pi_i)(y) = 1$. Hence it is clear that (X, t) is $\alpha - R_0(i)$.

Conversely, suppose that (X, t) is $\alpha - R_0(i)$. We shall prove that (X_i, t_i) , $i \in \Lambda$, is $\alpha - R_0(i)$. Let for some $i \in \Lambda$, a_i be a fixed element in X_i , suppose that $A_i = \{x \in X = \prod_{i \in \Lambda} X_i / x_j = a_j \text{ for some } i \neq j\}$. So that A_i is the subset of X , and this implies that (A_i, t_{A_i}) is also the subspace of (X, t) . Since (X, t) is $\alpha - R_0(i)$, then (A_i, t_{A_i}) is also $\alpha - R_0(i)$ and A_i is a homeomorphic image of X_i . Hence it is clear that (X_i, t_i) is $\alpha - R_0(i)$. ie (a) is proved.

Similarly (b), (c) and (d) can be proved.

2.2.1. Definition :-


Let (X, τ) be a fuzzy topological space and $\alpha \in I_1$.

(a) (X, τ) is said to be $\alpha - R_1(i) \Leftrightarrow \forall x, y \in X$ with $x \neq y$, whenever $\exists w \in \tau$ with $w(x) \neq w(y)$, then $\exists u, v \in \tau$ such that $u(x) = v(y) = 1$ and $u \cap v \leq \alpha$.

(b) (X, τ) is said to be $\alpha - R_1(ii) \Leftrightarrow \forall x, y \in X$ with $x \neq y$, whenever $\exists w \in \tau$ with $w(x) \neq w(y)$, then $\exists u, v \in \tau$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$.

(c) (X, τ) is said to be $\alpha - R_1(iii) \Leftrightarrow \forall x, y \in X$ with $x \neq y$, whenever $\exists w \in \tau$ with $w(x) \neq w(y)$, then $\exists u, v \in \tau$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$.

2.2.2. Lemma :- The following implications are true :



Proof :- First, suppose that (X, τ) is $\alpha - R_1(i)$. We shall prove that (X, τ) is $\alpha - R_1(iii)$.

Let $x, y \in X$ with $x \neq y$ and $w \in \tau$ such that $w(x) \neq w(y)$. Since (X, τ) is $\alpha - R_1(i)$, for $\alpha \in I_1, \exists u, v \in \tau$ such that $u(x) = v(y) = 1$ and $u \cap v \leq \alpha$. Now it is clear that when $w \in \tau$ with $w(x) \neq w(y), \exists u, v \in \tau$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. Hence it is clear that (X, τ) is $\alpha - R_1(iii)$.

Next, suppose that (X, τ) is $\alpha - R_1(ii)$. We shall prove that (X, τ) is $\alpha - R_1(iii)$. Let $x, y \in X$ with $x \neq y$ and $w \in \tau$ such that $w(x) \neq w(y)$. Since (X, τ) is $\alpha - R_1(ii)$, for $\alpha \in I_1, \exists u, v \in \tau$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$, ie $u \cap v \leq \alpha$. Hence it is clear that (X, τ) is $\alpha - R_1(iii)$.

Now, we give some examples to show the non implications among $\alpha - R_1(i)$, $\alpha - R_1(ii)$ and $\alpha - R_1(iii)$.

2.2.3. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1$, $u(y) = 0.4$, $v(x) = 0.4$, $v(y) = 1$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha > 0.4$, we see that (X, t) is $\alpha - R_1(i)$, but (X, t) is not $\alpha - R_1(ii)$.

2.2.4. Example :- Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.9$, $v(y) = 0.9$ and $v(x) = 0$, $u(y) = 0$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For every $\alpha < 0.9$. It is seen that (X, t) is $\alpha - R_1(ii)$, but (X, t) is not $\alpha - R_1(i)$.

2.2.5. Example: - Let $X = \{ x, y \}$, and $u, v \in I^X$, where u, v are defined by $u(x) = 0.8$, $v(y) = 0.8$ and $u(y) = 0.2$, $v(x) = 0.1$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For every $0.2 < \alpha < 0.8$, we have that (X, t) is $\alpha - R_1(iii)$, but (X, t) is not $\alpha - R_1(i)$ and (X, t) is not $\alpha - R_1(ii)$.

This completes the proof.

2.2.6. Theorem :- If $0 \leq \alpha \leq \beta < 1$ then

- (a) (X, t) is $\alpha - R_1(i)$. $\Rightarrow (X, t)$ is $\beta - R_1(i)$.
- (b) (X, t) is $\beta - R_1(ii)$. $\Rightarrow (X, t)$ is $\alpha - R_1(ii)$.
- (c) (X, t) is $0 - R_1(ii)$. $\Leftrightarrow (X, t)$ is $0 - R_1(iii)$.

Proof :- First, suppose that (X, t) is a $\alpha - R_1(i)$ space. We shall prove that (X, t) is a $\beta - R_1(i)$ space. Let $x, y \in X$ with $x \neq y$ and $w \in t$ such that $w(x) \neq w(y)$. Since (X, t) is $\alpha - R_1(i)$, for some $\alpha \in I_1$, $\exists u, v \in t$, such that $u(x) = 1 = v(y)$ and $u \cap v \leq \alpha$.

Since $0 \leq \alpha \leq \beta < 1$. Now we have $u \cap v \leq \beta$. Hence it is clear that (X, t) is a $\beta - R_1(i)$ space.

3.2.7. Example : - Let $X = \{x, y\}$, and $u, v \in I^X$, where u, v are defined by $u(x)=1, v(y)=1$ and $u(y)=0.2, v(x)=0.2$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.2$ and $\beta = 0.5$. It is clear that (X, t) is $\beta - R_1(i)$, but (X, t) is not $\alpha - R_1(i)$.

Next, suppose that (X, t) is $\beta - R_1(ii)$. We shall prove that (X, t) is $\alpha - R_1(ii)$. Let $x, y \in X$ with $x \neq y$ and $w \in t$ such that $w(x) \neq w(y)$. Since (X, t) is $\beta - R_1(ii)$, for $\beta \in I_1$, $\exists u, v \in t$ such that $u(x) > \beta, v(y) > \beta$ and $u \cap v = 0$. This implies that $u(x) > \alpha, v(y) > \alpha$ as $0 \leq \alpha \leq \beta < 1$. Hence it is clear that (X, t) is $\alpha - R_1(ii)$.

3.2.8. Example : - Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.6, v(y) = 0.6, u(y) = 0, v(x) = 0$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$ and $\beta = 0.8$. It is clear that (X, t) is $\alpha - R_1(ii)$, but (X, t) is not $\beta - R_1(ii)$.

Further, suppose that (X, t) is $0 - R_1(ii)$. We shall prove that (X, t) is $0 - R_1(iii)$. Let $x, y \in X$ with $x \neq y$ and $w \in t$ such that $w(x) \neq w(y)$. Since (X, t) is $0 - R_1(ii)$, $\exists u, v \in t$ such that $u(x) > 0, v(y) > 0$ and $u \cap v = 0$, ie $u \cap v \leq 0$. Hence it is clear that (X, t) is $0 - R_1(iii)$.

Conversely, suppose that (X, t) is $0 - R_1(iii)$. We shall prove that (X, t) is $0 - R_1(ii)$. Let $x, y \in X$ with $x \neq y$ and $w \in t$ such that $w(x) \neq w(y)$. Since (X, t) is $0 - R_1(iii)$, $\exists u, v \in t$ such that $u(x) > 0, v(y) > 0, u \cap v \leq 0, u \cap v = 0$. Hence (X, t) is $0 - R_1(ii)$. This completes the proof.

2.2.9. Theorem :- Let (X, t) be a fuzzy topological space, and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$ then

(a) (X, t) is an $\alpha - R_1$ (i) $\Rightarrow (X, I_\alpha(t))$ is a R_1 .

(b) (X, t) is an $\alpha - R_1$ (ii) $\Rightarrow (X, I_\alpha(t))$ is a R_1 .

(c) (X, t) is an $\alpha - R_1$ (iii) $\Rightarrow (X, I_\alpha(t))$ is a R_1 .

Proof :- Let (X, t) be $\alpha - R_1$ (i) space. We shall prove that $(X, I_\alpha(t))$ is R_1 - space. Let $x, y \in X$, with $x \neq y$ and $M \in I_\alpha(t)$ such that $x \in M, y \notin M$. We can write $M = w^{-1}(\alpha, 1]$, where $w \in t$. Then we see that $w(x) > \alpha, w(y) \leq \alpha$, therefore $w(x) \neq w(y)$. Since (X, t) is $\alpha - R_1$, for $\alpha \in I_1, \exists u, v \in t$, such that $u(x) = 1 = v(y)$ and $u \cap v \leq \alpha$. It follows that $\exists u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$ and $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ as $u \cap v \leq \alpha$. Thus it is clear that $(X, I_\alpha(t))$ is R_1 .

Next, suppose that (X, t) is $\alpha - R_1$ (ii). We shall prove that $(X, I_\alpha(t))$ is R_1 - space. Let $x, y \in X$ with $x \neq y$ and $M \in I_\alpha(t)$ such that $x \in M, y \notin M$. So we can write $M = w^{-1}(\alpha, 1]$ where $w \in t$. Then we see that $w(x) > \alpha, w(y) \leq \alpha$, and hence $w(x) \neq w(y)$. Since (X, t) is $\alpha - R_1$ (ii), for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$. It follows that $\exists u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$ and $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ as $u(x) > \alpha, v(y) > \alpha$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$ as $u \cap v = 0$. Now it is clear that $(X, I_\alpha(t))$ is R_1 - Space.

Finally, Suppose that (X, t) is $\alpha - R_1$ (iii) space. We shall prove that $(X, I_\alpha(t))$ is a R_1 space. Let $x, y \in X$, with $x \neq y$ and $M \in I_\alpha(t)$ such that $x \in M, y \notin M$ or $x \notin M, y \in M$. Suppose that $x \in M, y \notin M$. We can write $M = w^{-1}(\alpha, 1]$, where $w \in t$. Now we have $w(x) > \alpha, w(y) \leq \alpha$ ie $w(x) \neq w(y)$. Since (X, t) is $\alpha - R_1$ (iii), for $\alpha \in I_1$,

then $\exists u, v \in \tau$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. It follows that $\exists u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(\tau)$ with $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ as $u(x) > \alpha, v(y) > \alpha$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$, since $u \cap v \leq \alpha$. Now it is clear that (X, τ) is R_1 .

Now we give some examples.

2.2.10. Example : - Let $X = \{x, y\}$ and $u, v, w \in I^X$, where u, v and w are defined by $u(x) = 1, u(y) = 0.1, v(x) = 0.4, v(y) = 0.9, w(x) = 0.7, w(y) = 0.3$. Consider the fuzzy topology τ on X generated by $\{0, u, v, w, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$, it is clear that (X, τ) is not $\alpha - R_1(i)$ and (X, τ) is not $\alpha - R_1(ii)$. Now $I_\alpha(\tau) = \{X, \phi, \{x\}, \{y\}\}$. Then clear that $I_\alpha(\tau)$ is a topology on X and $(X, I_\alpha(\tau))$ is R_1 -Space.


2.2.11. Example:- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.4, u(y) = 0.5$ and $v(x) = 0.3, v(y) = 0.4$. Consider the fuzzy topology τ on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$, we see that (X, τ) is not $\alpha - R_1(iii)$. Now $I_\alpha(\tau) = \{X, \phi\}$. Then $I_\alpha(\tau)$ is a topology on X and $(X, I_\alpha(\tau))$ is R_1 -Space.

This completes the proof.

2.2.12. Theorem :- Let (X, T) be a topological space. Consider the following statements:

- (1) (X, T) be a R_1 space.
- (2) $(X, \omega(T))$ be an $\alpha - R_1(i)$ space.
- (3) $(X, \omega(T))$ be an $\alpha - R_1(ii)$ space.
- (4) $(X, \omega(T))$ be an $\alpha - R_1(iii)$ space.

Then the following implications are true.



Proof :- First suppose that (X, T) be a R_1 -space . We shall prove that $(X, \omega(T))$ be a α - R_1 (i) space. Let $x, y \in X$ with $x \neq y$ and $m \in \omega(T)$ such that $m(x) \neq m(y)$ ie either $m(x) < m(y)$ or $m(x) > m(y)$. Suppose that $m(x) < r < m(y)$. Then it is clear that $m^{-1}(r, 1] \in T$ as $m \in \omega(T)$ and $x \notin m^{-1}(r, 1], y \in m^{-1}(r, 1]$. Since (X, T) is R_1 space then $\exists U, V \in T$ such that $x \in U, y \in V$ and $U \cap V = \phi$. Since $1_U, 1_V$ are lower semi continuous function from (X, T) into I , then $1_U, 1_V \in \omega(T)$, and it is clear that $1_U(x) = 1, 1_V(y) = 1$ and $1_U \cap 1_V = 0$ ie $(X, \omega(T))$ is α - R_1 (i) space . Also it is clear that $(X, \omega(T))$ is α - R_1 (ii) space .

Further, it is easy to show that $(2) \Rightarrow (4)$ and $(3) \Rightarrow (4)$.

We, therefore prove that $(4) \Rightarrow (1)$.

Suppose that $(X, \omega(T))$ is α - R_1 (iii). We shall prove that (X, T) is R_1 space . Let $x, y \in X$ with $x \neq y$ and $M \in T$ such that $x \in M, y \notin M$ or $x \notin M, y \in M$. Suppose $x \in M, y \notin M$. But 1_M is lower semi continuous function from (X, T) into I , so $1_M \in \omega(T)$ and $1_M(x) = 1, 1_M(y) = 0$ ie $1_M(x) \neq 1_M(y)$. Since $(X, \omega(T))$ is α - R_1 (iii), for $\alpha \in I_1$ then $\exists u, v \in \omega(T)$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. Now we is observed that $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in T$ such that, $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$. Thus (X, T) is R_1 space .

This completes the proof.

Thus it is seen that $\alpha - R_1(p)$ is a good extension of its topological counter part . ($p = i, ii, iii,$)

2.2.13. Theorem :- Let (X, t) be a fuzzy topological space, $A \subseteq X$ and $t_A = \{ u/A : u \in t \}$, then

- (a) (X, t) is $\alpha - R_1(i) \Rightarrow (A, t_A)$ is $\alpha - R_1(i)$.
- (b) (X, t) is $\alpha - R_1(ii) \Rightarrow (A, t_A)$ is $\alpha - R_1(ii)$.
- (c) (X, t) is $\alpha - R_1(iii) \Rightarrow (A, t_A)$ is $\alpha - R_1(iii)$.

Proof :- Suppose (X, t) is $\alpha - R_1(iii)$. We shall prove that (A, t_A) is $\alpha - R_1(iii)$. Let $x, y \in A$ with $x \neq y$ then $x, y \in X$ and $x \neq y$. Consider $m \in t_A$ with $m(x) \neq m(y)$. Then m can be written as w/A , where $w \in t$ and hence $w(x) \neq w(y)$. Since (X, t) is $\alpha - R_1(iii)$, for $\alpha \in I_1$, then $\exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. But we have $u/A \in t_A$, for every $u \in t$. Now we observed that $u/A(x) > \alpha, v/A(y) > \alpha$ and $u/A \cap v/A \leq \alpha$, since $u \cap v \leq \alpha$. Hence it is clear that (A, t_A) is $\alpha - R_1(iii)$.

Similarly (a) and (b) can be proved .

2.2.14. Theorem :- Given $(X_i, t_i), i \in \Lambda$ be fuzzy topological spaces and $X = \prod_{i \in \Lambda} X_i$ and t be the product topology on X , then

- (a) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - R_1(i) \Leftrightarrow (X, t)$ is $\alpha - R_1(i)$.
- (b) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - R_1(ii) \Leftrightarrow (X, t)$ is $\alpha - R_1(ii)$.
- (c) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - R_1(iii) \Leftrightarrow (X, t)$ is $\alpha - R_1(iii)$.

Proof :- Suppose that $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - R_1(iii)$. We shall prove that (X, t) is $\alpha - R_1(iii)$. Let $x, y \in X$ with $x \neq y$ and $w \in t$ with $w(x) \neq w(y)$. But we have $w(x) = \min \{ w_i(x_i) : i \in \Lambda \}, w(y) = \min \{ w_i(y_i) : i \in \Lambda \}$. Hence we can find at least one

$w_i \in t_i$ and $x_i, y_i \in X_i$, with $x_i \neq y_i$ and $w_i(x_i) \neq w_i(y_i)$. Since $(X_i, t_i), i \in \Lambda$ is α - R_1 (iii), for $\alpha \in I_1$, then $\exists u_i, v_i \in t_i$ such that $u_i(x_i) > \alpha$, $v_i(y_i) > \alpha$ and $u_i \cap v_i \leq \alpha$. But we have $\pi_i(x) = x_i$ and $\pi_i(y) = y_i$ and hence $u_i(\pi_i(x)) > \alpha$, $v_i(\pi_i(y)) > \alpha$. It follows that $\exists u_i \circ \pi_i, v_i \circ \pi_i \in t$ such that $(u_i \circ \pi_i)(x) > \alpha$, $(v_i \circ \pi_i)(y) > \alpha$ and $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) \leq \alpha$. Hence it is clear that (X, t) is α - R_1 (iii).

Conversely, suppose that (X, t) is α - R_1 (iii). We shall prove that $(X_i, t_i), i \in \Lambda$, is α - R_1 (iii). Let for some $i \in \Lambda$, a_i be a fixed element in X_i , suppose that $A_i = \{x \in X = \prod_{i \in \Lambda} X_i / x_j = a_j \text{ for some } i \neq j\}$. So that A_i is a subset of X , and hence (A_i, t_{A_i}) is also a subspace of (X, t) . Since (X, t) is α - R_1 (iii), then (A_i, t_{A_i}) is also α - R_1 (iii). Now we have A_i is homeomorphic image of X_i . Hence it is clear that (X_i, t_i) is α - R_1 (iii) ie (c) is proved.

Similarly (a) and (b) can be proved.

This completes the proof.

Chapter 3

T_0 and T_1 Fuzzy Topological spaces

3.1 Introduction :-

In this chapter , we introduce and study some T_0 and T_1 properties in fuzzy topological spaces and obtain their several features .

3.1.1. Definition :-

Let (X, τ) be a fuzzy topological space and $\alpha \in I_1$.


(a) (X, τ) is an $\alpha - T_0(i)$ space $\Leftrightarrow \forall x, y \in X$, with $x \neq y$, $\exists u \in \tau$ such that $u(x) = 1$, $u(y) \leq \alpha$ or $\exists v \in \tau$ such that $v(x) \leq \alpha$, $v(y) = 1$.

(b) (X, τ) is an $\alpha - T_0(ii)$ space $\Leftrightarrow \forall x, y \in X$, with $x \neq y$, $\exists u \in \tau$ such that $u(x) = 0$, $u(y) > \alpha$ or $\exists v \in \tau$ such that $v(x) > \alpha$, $v(y) = 0$.

(c) (X, τ) is an $\alpha - T_0(iii)$ space $\Leftrightarrow \forall x, y \in X$, with $x \neq y$, $\exists u \in \tau$ such that $0 \leq u(x) \leq \alpha < u(y) \leq 1$ or $\exists v \in \tau$ such that $0 \leq v(y) \leq \alpha < v(x) \leq 1$.

(d) (X, τ) is a $T_0(iv)$ space $\Leftrightarrow \forall x, y \in X$, with $x \neq y$, $\exists u \in \tau$ such that $u(x) \neq u(y)$.

3.1.2. Lemma: - The following implications are true :



Proof : - Let (X, τ) be a fuzzy topological space and (X, τ) is $\alpha - T_0$ (i) . We shall prove that (X, τ) is $\alpha - T_0$ (iii). Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $\alpha - T_0$ (i) , fore $\alpha \in I_1$, $\exists u \in \tau$ such that $u(x)=1, u(y) \leq \alpha$. It follows that $0 \leq u(y) \leq \alpha < u(x) \leq 1$. Hence it is clear that (X, τ) is $\alpha - T_0$ (iii) .

Next, suppose that (X, τ) is $\alpha - T_0$ (ii) . We shall prove that (X, τ) is $\alpha - T_0$ (iii). Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $\alpha - T_0$ (ii) , for $\alpha \in I_1$, $\exists u \in \tau$ such that $u(x) = 0, u(y) > \alpha$. This implies that $0 \leq u(x) \leq \alpha < u(y) \leq 1$. Hence it is clear that (X, τ) is $\alpha - T_0$ (iii).

Finally, suppose that (X, τ) is $\alpha - T_0$ (iii) . We shall prove that (X, τ) is T_0 (iv) . Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $\alpha - T_0$ (iii) , for $\alpha \in I_1$, $\exists u \in \tau$ such that $0 \leq u(x) \leq \alpha < u(y) \leq 1$. Now we observe that $u(x) \neq u(y)$. Hence (X, τ) is T_0 (iv) .

Now , we give some examples to show the nonimplications among $\alpha - T_0$ (i) , $\alpha - T_0$ (ii), $\alpha - T_0$ (iii) and T_0 (iv) .

3.1.3. Example: - Let $X = \{ x, y \}$ and $u \in I^X$ be given by $u(x) = 0.6, u(y) = 0.8$. Consider the fuzzy topology τ on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.7$ it is clear that (X, τ) is $\alpha - T_0$ (iii) but (X, τ) is not $\alpha - T_0$ (i) . It is also clear that (X, τ) is not $\alpha - T_0$ (ii).

3.1.4. Example : - Let $X = \{ x, y \}$ and $u \in I^X$, where u is defined by $u(x) = 1, u(y) = 0.4$. Consider the fuzzy topology τ on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$, we see that (X, τ) is $\alpha - T_0$ (i) but (X, τ) is not $\alpha - T_0$ (ii) .

3.1.5. Example : - Let $X = \{ x, y \}$ and $u \in I^X$, where u is defined by $u(x) = 0, u(y) = 0.7$. Consider the fuzzy topology τ on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.4$, it is clear that (X, τ) is $\alpha - T_0$ (ii) but (X, τ) is not $\alpha - T_0$ (i) .

3.1.6. Example : - Let $X = \{ x, y \}$ and $u \in I^X$, where u is defined by $u(x) = 0.4$, $u(y) = 0.8$. Consider the fuzzy topology t on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.9$, we see that (X, t) is T_0 (iv) but (X, t) is not $\alpha - T_0$ (iii).

This completes the proof.

3.1.7. Lemma :- If $0 \leq \alpha \leq \beta < 1$, then

(a) (X, t) is $\alpha - T_0$ (i) $\Rightarrow (X, t)$ is $\beta - T_0$ (i).

(b) (X, t) is $\beta - T_0$ (ii) $\Rightarrow (X, t)$ is $\alpha - T_0$ (ii).

(c) (X, t) is $0 - T_0$ (ii) $\Leftrightarrow (X, t)$ is $0 - T_0$ (iii).

Proof :- Suppose that (X, t) be a fuzzy topological space and (X, t) is $\alpha - T_0$ (i). We shall prove that (X, t) is $\beta - T_0$ (i). Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_0$ (i), for $\alpha \in I_1$, $\exists u \in t$ such that $u(x) = 1$, $u(y) \leq \alpha$. This implies that $u(x) = 1$, $u(y) \leq \beta$, since $0 \leq \alpha \leq \beta < 1$. Hence it is clear that (X, t) is $\beta - T_0$ (i).

3.1.8. Example: - Let $X = \{ x, y \}$ and $u \in I^X$, where u is defined by $u(x) = 1$, $u(y) = 0.7$. Consider the fuzzy topology t on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$ and $\beta = 0.8$, we see that (X, t) is $\beta - T_0$ (i) but (X, t) is not $\alpha - T_0$ (i).

Next, suppose that (X, t) is $\beta - T_0$ (ii). We shall prove that (X, t) is $\alpha - T_0$ (ii).

Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\beta - T_0$ (ii), for $\beta \in I_1$, $\exists u \in t$ such that $u(x) = 0$, $u(y) > \beta$. This implies that $u(x) = 0$, $u(y) > \alpha$, since $0 \leq \alpha \leq \beta < 1$. Hence it is clear that (X, t) is $\alpha - T_0$ (ii).

3.1.9. Example:- Let $X = \{ x, y \}$ and $u \in I^X$, where u is defined by $u(x) = 0$, $u(y) = 0.5$. Consider the fuzzy topology t on X generated by $\{ 0, u, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.4$ and $\beta = 0.7$, we see that (X, t) is $\alpha - T_0$ (ii), but (X, t) is not $\beta - T_0$ (ii).

Finally, suppose that (X, τ) is $0-T_0$ (ii). We shall prove that (X, τ) is $0-T_0$ (iii).

Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $0-T_0$ (ii), $\exists u \in \tau$ such that $u(x) = 0, u(y) > 0$.

This implies that $0 \leq u(x) \leq 0 < u(y) \leq 1$. Hence (X, τ) is $0-T_0$ (iii).

Conversely, suppose that (X, τ) is $0-T_0$ (iii). We shall prove that (X, τ) is $0-T_0$ (ii).


Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $0-T_0$ (iii), $\exists u \in \tau$ such that $0 \leq u(x) \leq 0 < u(y) \leq 1$ ie $u(x) = 0, u(y) > 0$. Hence (X, τ) is $0-T_0$ (ii).

This completes the proof.

3.1.10. Theorem :- Let (X, T) be a topological space. Consider the following statements:

- (1) (X, T) be a T_0 - space.
- (2) $(X, \omega(T))$ be an $\alpha-T_0$ (i) space .
- (3) $(X, \omega(T))$ be an $\alpha-T_0$ (ii) space .
- (4) $(X, \omega(T))$ be an $\alpha-T_0$ (iii) space .
- (5) $(X, \omega(T))$ be a T_0 (iv) space .

Then the implications given below are true :



Proof :- Suppose (X, T) is a T_0 - topological space . We shall prove that $(X, \omega(T))$ is $\alpha-T_0$ (i) fuzzy topological space. Let $x, y \in X$ with $x \neq y$. Since (X, T) is T_0 , $\exists U \in T$ such that $x \in U, y \notin U$. But from the definition of lsc , we have $1_U \in \omega(T)$ and $1_U(x) = 1, 1_U(y) = 0$. Hence we see that $(X, \omega(T))$ is $\alpha-T_0$ (i) space . Also we see that $(X, \omega(T))$ is $\alpha-T_0$ (ii) space.

Further, it is easy to show that (2) \Rightarrow (4), (3) \Rightarrow (4) and (4) \Rightarrow (5).

We, therefore prove that (4) \Rightarrow (1).

Suppose $(X, \omega(T))$ be a $T_0(iv)$ space. We shall prove that (X, T) is T_0 - space. Let $x, y \in X$ with $x \neq y$. Since $(X, \omega(T))$ is $T_0(iv)$, $\exists u \in \omega(T)$ such that $u(x) \neq u(y)$ ie either $u(x) < u(y)$ or $u(x) > u(y)$. Suppose $u(x) < u(y)$. Then we can find a $r \in I_1$, such that $u(x) < r < u(y)$. We observe that $x \notin u^{-1}(r, 1]$, $y \in u^{-1}(r, 1]$, and by the definition of lsc , $u^{-1}(r, 1] \in T$. Hence (X, T) is T_0 - space.

Thus it is seen that $\alpha - T_0(p)$ is a good extension of its topological counter part ($p = i, ii, iii, iv$).

This completes the proof.

3.1.11. Theorem : - Let (X, t) be a fuzzy topological space, and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$, then

- (a) (X, t) is an $\alpha - T_0(i) \Rightarrow (X, I_\alpha(t))$ is T_0 .
- (b) (X, t) is an $\alpha - T_0(ii) \Rightarrow (X, I_\alpha(t))$ is T_0 .
- (c) (X, t) is an $\alpha - T_0(iii) \Leftrightarrow (X, I_\alpha(t))$ is T_0 .

Proof :- (a) Let (X, t) be a fuzzy topological space and (X, t) be $\alpha - T_0(i)$. We shall prove that $(X, I_\alpha(t))$ is T_0 - space. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_0(i)$, fore $\alpha \in I_1$, $\exists u \in t$ such that $u(x) = 1, u(y) \leq \alpha$. Since $u^{-1}(\alpha, 1] \in I_\alpha(t)$, $y \notin u^{-1}(\alpha, 1]$ and $x \in u^{-1}(\alpha, 1]$. We have that $(X, I_\alpha(t))$ is T_0 - space.

(b) Again, suppose that (X, t) is $\alpha - T_0(ii)$. We shall prove that $(X, I_\alpha(t))$ is T_0 - space. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_0(ii)$, for $\alpha \in I_1$, $\exists u \in t$ such that $u(x) = 0$,

$u(y) > \alpha$. Since $u^{-1}(\alpha, 1] \in I_\alpha(t)$, $x \notin u^{-1}(\alpha, 1]$ and $y \in u^{-1}(\alpha, 1]$, one can see that $(X, I_\alpha(t))$ is T_0 -space.

(c) Finally, suppose that (X, t) is α - T_0 (iii). We shall prove that $(X, I_\alpha(t))$ is T_0 -space. Let $x, y \in X$ with $x \neq y$. Since (X, t) is α - T_0 (iii), for $\alpha \in I_1$, $\exists u \in t$ such that $0 \leq u(x) \leq \alpha < u(y) \leq 1$. Since $u^{-1}(\alpha, 1] \in I_\alpha(t)$, $x \notin u^{-1}(\alpha, 1]$ and $y \in u^{-1}(\alpha, 1]$, so it is clear that $(X, I_\alpha(t))$ is T_0 -space.

Conversely, suppose $(X, I_\alpha(t))$ be T_0 -space. We shall prove that (X, t) is a α - T_0 (iii) space. Let $x, y \in X$ with $x \neq y$. Since $(X, I_\alpha(t))$ is T_0 -space, $\exists u^{-1}(\alpha, 1] \in I_\alpha(t)$ such that $x \in u^{-1}(\alpha, 1]$ and $y \notin u^{-1}(\alpha, 1]$, where $u \in t$. Thus we have that $u(x) > \alpha$, $u(y) \leq \alpha$, ie $0 \leq u(y) \leq \alpha < u(x) \leq 1$, one can see that (X, t) is α - T_0 (iii) space.

Now we give an example:-

3.1.12. Example :- Let $X = \{x, y\}$ and $u \in I^X$, where u is defined by $u(x) = 0.8$, $u(y) = 0.2$.

Consider the fuzzy topology t on X generated by $\{0, u, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$ it is clear that (X, t) is not α - T_0 (i) and (X, t) is not α - T_0 (ii). Now $I_\alpha(t) = \{X, \phi, \{x\}\}$.

Then we see that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is a T_0 space.

This completes the proof.

3.1.13. Theorem :- Let (X, t) be a fuzzy topological space, $A \subseteq X$, $t_A = \{u/A : u \in t\}$,

then,

- (a) (X, t) is α - T_0 (i) $\Rightarrow (A, t_A)$ is α - T_0 (i).
- (b) (X, t) is α - T_0 (ii) $\Rightarrow (A, t_A)$ is α - T_0 (ii).
- (c) (X, t) is α - T_0 (iii) $\Rightarrow (A, t_A)$ is α - T_0 (iii).
- (d) (X, t) is T_0 (iv) $\Rightarrow (A, t_A)$ is T_0 (iv).

Proof :- Suppose that (X, τ) be a fuzzy topological space and (X, τ) is $\alpha - T_0(i)$. We shall prove that (A, τ_A) is $\alpha - T_0(i)$. Let $x, y \in A$ with $x \neq y$, so that $x, y \in X$, as $A \subseteq X$. Since (X, τ) is $\alpha - T_0(i)$, for $\alpha \in I_1, \exists u \in \tau$ such that $u(x) = 1, u(y) \leq \alpha$. For $A \subseteq X$, we have $u|_A \in \tau_A$ and $u|_A(x) = 1, u|_A(y) \leq \alpha$ as $x, y \in A$. Hence it is clear that (A, τ_A) is $\alpha - T_0(i)$.

Similarly (b), (c) and (d) can be proved.

3.1.14. Theorem :- Given $(X_i, \tau_i), i \in \Lambda$ be fuzzy topological spaces and $X = \prod_{i \in \Lambda} X_i$ and τ be the product topology on X , then

- (a) $\forall i \in \Lambda, (X_i, \tau_i)$ is $\alpha - T_0(i) \Leftrightarrow (X, \tau)$ is $\alpha - T_0(i)$.
- (b) $\forall i \in \Lambda, (X_i, \tau_i)$ is $\alpha - T_0(ii) \Leftrightarrow (X, \tau)$ is $\alpha - T_0(ii)$.
- (c) $\forall i \in \Lambda, (X_i, \tau_i)$ is $\alpha - T_0(iii) \Leftrightarrow (X, \tau)$ is $\alpha - T_0(iii)$.
- (d) $\forall i \in \Lambda, (X_i, \tau_i)$ is $T_0(iv) \Leftrightarrow (X, \tau)$ is $T_0(iv)$.

Proof :- Suppose that $\forall i \in \Lambda, (X_i, \tau_i)$ is $\alpha - T_0(i)$. We shall prove that (X, τ) is $\alpha - T_0(i)$. Let $x, y \in X$ with $x \neq y$, then $x_i \neq y_i$, for some $i \in \Lambda$. Since (X_i, τ_i) is $\alpha - T_0(i)$, for $\alpha \in I_1, \exists u_i \in \tau_i, i \in \Lambda$ such that $u_i(x_i) = 1$ and $u_i(y_i) \leq \alpha$. But we have $\pi_i(x) = x_i$, and $\pi_i(y) = y_i$. Then $u_i(\pi_i(x)) = 1$ and $u_i(\pi_i(y)) \leq \alpha$, ie $(u_i \circ \pi_i)(x) = 1$ and $(u_i \circ \pi_i)(y) \leq \alpha$. It follows that $\exists (u_i \circ \pi_i) \in \tau$ such that $(u_i \circ \pi_i)(x) = 1, (u_i \circ \pi_i)(y) \leq \alpha$. Hence it is clear that (X, τ) is $\alpha - T_0(i)$.

Conversely, suppose that (X, τ) is $\alpha - T_0(i)$ space. We shall prove that $(X_i, \tau_i), i \in \Lambda$ is $\alpha - T_0(i)$. Let a_j be a fixed element in X_j , suppose that $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$. So that A_i is a subset of X , and hence (A_i, τ_{A_i}) is also a subspace of (X, τ) . Since (X, τ) is $\alpha - T_0(i)$ then (A_i, τ_{A_i}) is also $\alpha - T_0(i)$. Now we have A_i is homeomorphic image of X_i . Hence it is clear that $(X_i, \tau_i), i \in \Lambda$ is $\alpha - T_0(i)$.

Similarly (b) , (c) and (d) can be proved .

3.1.15. Theorem:- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be a one-one , onto and open map then,

$$(a) (X, t) \text{ is } \alpha\text{-}T_0(i) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_0(i) .$$

$$(b) (X, t) \text{ is } \alpha\text{-}T_0(ii) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_0(ii) .$$

$$(c) (X, t) \text{ is } \alpha\text{-}T_0(iii) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_0(iii) .$$

$$(d) (X, t) \text{ is } T_0(iv) \Rightarrow (Y, s) \text{ is } T_0(iv) .$$

Proof :- Suppose (X, t) be $\alpha\text{-}T_0(i)$. We shall prove that (Y, s) is $\alpha\text{-}T_0(i)$. Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto then , $\exists x_1, x_2 \in X$ with $f(x_1) = y_1$, $f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one. Again since (X, t) is $\alpha\text{-}T_0(i)$, for $\alpha \in I_1$, then $\exists u \in t$ such that $u(x) = 1$, $u(y) \leq \alpha$.

$$\text{Now } f(u)(y_1) = \{ \text{Sup } u(x_1) : f(x_1) = y_1 \}$$

$$= 1 .$$

$$f(u)(y_2) = \{ \text{Sup } u(x_2) : f(x_2) = y_2 \}$$

$$\leq \alpha .$$

Since f is open then $f(u) \in s$ as $u \in t$. We observe that $\exists f(u) \in s$ such that $f(u)(y_1) = 1$,

$f(u)(y_2) \leq \alpha$. Hence it is clear that (Y, s) is $\alpha\text{-}T_0(i)$.

Similarly (b) , (c) and (d) can be proved .

3.1.16. Theorem :- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be continuous and one-one map, then

$$(a) (Y, s) \text{ is } \alpha\text{-}T_0(i) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_0(i) .$$

$$(b) (Y, s) \text{ is } \alpha\text{-}T_0(ii) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_0(ii) .$$

$$(c) (Y, s) \text{ is } \alpha\text{-}T_0(iii) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_0(iii) .$$

(d) (Y, s) is $T_0(iv) \Rightarrow (X, t)$ is $T_0(iv)$.

Proof:- Suppose (Y, s) be $\alpha - T_0(i)$. We shall prove that (X, t) is $\alpha - T_0(i)$. Let $x_1, x_2 \in X$ with $x_1 \neq x_2$, then $f(x_1) \neq f(x_2)$ in Y , as f is one-one. Since (Y, s) is $\alpha - T_0(i)$, for $\alpha \in I_1$, then $\exists u \in s$ such that $u(f(x_1)) = 1, u(f(x_2)) \leq \alpha$. This implies that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) \leq \alpha$, since $u \in s$ and f is continuous then $f^{-1}(u) \in t$. Now it is clear that $\exists f^{-1}(u) \in t$ such that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) \leq \alpha$. Hence (X, t) is $\alpha - T_0(i)$.
Similarly (b), (c) and (d) can be proved.

3.2.1. Definition :-

Let (X, t) be a fuzzy topological space and $\alpha \in I_1$.


(a) (X, t) is an $\alpha - T_1(i)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) = 1, u(y) \leq \alpha$ and $v(x) \leq \alpha, v(y) = 1$.

(b) (X, t) is an $\alpha - T_1(ii)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) = 0, u(y) > \alpha$ and $v(x) > \alpha, v(y) = 0$.

(c) (X, t) is an $\alpha - T_1(iii)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $0 \leq u(y) \leq \alpha < u(x) \leq 1$ and $0 \leq v(x) \leq \alpha < v(y) \leq 1$.

(d) (X, t) is a $T_1(iv)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) < u(y)$ and $v(x) > v(y)$.

3.2.2. Lemma :- The following implications are true:



Proof:- Let (X, t) be $\alpha\text{-}T_1(i)$ fuzzy topological space . We shall prove that (X, t) is $\alpha\text{-}T_1(iii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha\text{-}T_1(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 1, u(y) \leq \alpha$ and $v(x) \leq \alpha, v(y) = 1$. We see that $0 \leq u(y) \leq \alpha < u(x) \leq 1$ and $0 \leq v(x) \leq \alpha < v(y) \leq 1$. Hence it is clear that (X, t) is $\alpha\text{-}T_1(iii)$.

Next, suppose that (X, t) is $\alpha\text{-}T_1(ii)$. We shall prove that (X, t) is $\alpha\text{-}T_1(iii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha\text{-}T_1(ii)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 0, u(y) > \alpha$ and $v(x) > \alpha, v(y) = 0$. Now we see that $0 \leq u(x) \leq \alpha < u(y) \leq 1$ and $0 \leq v(y) \leq \alpha < v(x) \leq 1$. Hence (X, t) is $\alpha\text{-}T_1(iii)$.

Finally, suppose that (X, t) is $\alpha\text{-}T_1(iii)$. We shall prove that (X, t) is $T_1(iv)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha\text{-}T_1(iii)$, for $\alpha \in I_1, \exists u, v \in t$ such that $0 \leq u(x) \leq \alpha < u(y) \leq 1$ and $0 \leq v(y) \leq \alpha < v(x) \leq 1$. This implies that $u(x) < u(y)$ and $v(x) > v(y)$. Hence (x, t) is $T_1(iv)$.

Now we give some examples to show the non implication among $\alpha\text{-}T_1(i), \alpha\text{-}T_1(ii), \alpha\text{-}T_1(iii)$ and $T_1(iv)$.

3.2.3. Example:- Let $X = \{x, y\}$ and $u, v \in I^X$ where u, v are defined by $u(x) = 0.6, u(y) = 0.8$ and $v(x) = 0.8, v(y) = 0.6$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.7$, it is clear that (X, t) is $\alpha\text{-}T_1(iii)$. But (x, t) is not $\alpha\text{-}T_1(i)$ and (X, t) is not $\alpha\text{-}T_1(ii)$.

3.2.4. Example: - Let $X = \{ x, y \}$ and $u, v \in I^X$ where u, v are defined by $u(x) = 1$, $u(y) = 0.4$, and $v(x) = 0.4$, $v(y) = 1$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$, we see that (X, t) is α - $T_1(i)$, but (X, t) is not α - $T_1(ii)$.

3.2.5. Example : - Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0$, $u(y) = 0.7$ and $v(x) = 0.7$, $v(y) = 0$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.6$, we see that (X, t) is α - $T_1(ii)$, but (X, t) is not α - $T_1(i)$.

3.2.6. Example : - Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.4$, $u(y) = 0.6$ and $v(x) = 0.3$, $v(y) = 0.3$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.8$, we can see that (X, t) is $T_1(iv)$ but (X, t) is not α - $T_1(iii)$.

This completes the proof.

3.2.7. Theorem : - If $0 \leq \alpha \leq \beta < 1$, then

(a) (X, t) is α - $T_1(i) \Rightarrow (X, t)$ is β - $T_1(i)$.

(b) (X, t) is β - $T_1(ii) \Rightarrow (X, t)$ is α - $T_1(ii)$.

(c) (X, t) is 0 - $T_1(ii) \Leftrightarrow (X, t)$ is 0 - $T_1(iii)$.

Proof: - (a) Let (X, t) be α - $T_1(i)$ space. We shall prove that (X, t) is β - $T_1(i)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is α - $T_1(i)$, for $\alpha \in I_1$, $\exists u, v \in t$ such that $u(x) = 1$, $u(y) \leq \alpha$, and $v(x) \leq \alpha$, $v(y) = 1$. This implies that $u(x) = 1$, $u(y) \leq \beta$ and $v(x) \leq \beta$, $v(y) = 1$ as $0 \leq \alpha \leq \beta < 1$. Hence (X, t) is β - $T_1(i)$.

3.2.8. Example: - Let $X = \{ x, y \}$ and $u, v \in I^X$ where u, v are defined by $u(x) = 1$, $u(y) = 0.7$ and $v(x) = 0.7$, $v(y) = 1$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.5$ and $\beta = 0.8$, it is clear that (X, t) is $\beta - T_1(i)$ but (X, t) is not $\alpha - T_1(i)$.

(b) Next, suppose that (X, t) is $\beta - T_1(ii)$. We shall prove that (X, t) is $\alpha - T_1(ii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\beta - T_1(ii)$, for $\beta \in I_1$, $\exists u, v \in t$ such that $u(x) = 0$, $u(y) > \beta$ and $v(x) > \beta$, $v(y) = 0$. So, we see that $u(x) = 0$, $u(y) > \alpha$ and $v(x) > \alpha$, $v(y) = 0$ as $0 \leq \alpha \leq \beta < 1$. Hence (X, t) is $\alpha - T_1(ii)$.

3.2.9. Example : - Let $X = \{ x, y \}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0$, $u(y) = 0.5$, and $v(x) = 0.5$, $v(y) = 0$. Consider the fuzzy topology t on X generated by $\{ 0, u, v, 1 \} \cup \{ \text{Constants} \}$. For $\alpha = 0.3$ and $\beta = 0.7$. It is clear that (X, t) is $\alpha - T_1(ii)$ but (X, t) is not $\beta - T_1(ii)$.

(c) Finally, suppose (X, t) be $0 - T_1(ii)$. We shall prove that (X, t) is $0 - T_1(iii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $0 - T_1(ii)$, $\exists u, v \in t$ such that $u(x) = 0$, $u(y) > 0$ and $v(x) > 0$, $v(y) = 0$. So, we see that $0 \leq u(x) \leq 0 < u(y) \leq 1$ and $0 \leq v(y) \leq 0 < v(x) \leq 1$. Hence (X, t) is $0 - T_1(iii)$.


Conversely, suppose that (X, t) is $0 - T_1(iii)$. We shall prove that (X, t) is $0 - T_1(ii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $0 - T_1(iii)$, $\exists u, v \in t$ such that $0 \leq u(x) \leq 0 < u(y) \leq 1$ and $0 \leq v(y) \leq 0 < v(x) \leq 1$. Thus we see that $u(x) = 0$, $u(y) > 0$ and $v(y) = 0$, $v(x) > 0$. Hence (X, t) is $0 - T_1(ii)$.

This completes the proof.

3.2.10. Theorem: - Let (X, T) be a topological space. Consider the following statements.

- (1) (X, T) be a T_1 - space .
- (2) $(X, \omega(T))$ be an $\alpha - T_1(i)$ space .
- (3) $(X, \omega(T))$ be an $\alpha - T_1(ii)$ space .
- (4) $(X, \omega(T))$ be an $\alpha - T_1(iii)$ space .
- (5) $(X, \omega(T))$ be an $T_1(iv)$ space .

Then the implications are true:



Proof :- Let (X, T) be a T_1 space. We shall prove that $(X, \omega(T))$ is $\alpha - T_1(i)$. Let $x, y \in X$ with $x \neq y$. Since (X, T) is T_1 - space, $\exists U, V \in T$ such that $x \in U, y \notin U$ and $x \notin V, y \in V$. But from the definition of lsc, $1_U, 1_V \in \omega(T)$ and $1_U(x) = 1, 1_U(y) = 0$ and $1_V(x) = 0, 1_V(y) = 1$. Hence $(X, \omega(T))$ is $\alpha - T_1(i)$ space, for any $\alpha \in I_1$. Also it is clear that $(X, \omega(T))$ is $\alpha - T_1(ii)$.

Further, it is easy to show that $(2) \Rightarrow (4), (3) \Rightarrow (4)$ and $(4) \Rightarrow (5)$.

We, therefore prove that $(5) \Rightarrow (1)$.

Suppose, $(X, \omega(T))$ be $T_1(iv)$ space. We shall prove that (X, T) is T_1 - space. Let $x, y \in X$ with $x \neq y$. Since $(X, \omega(T))$ is $T_1(iv)$, $\exists u, v \in \omega(T)$ such that $u(x) < u(y)$ and $v(x) > v(y)$. Let $r, s \in I$ be such that $u(x) < r < u(y)$ and $v(x) > s > v(y)$. Then we have

$u^{-1}(r, 1], v^{-1}(s, 1] \in T$ and $x \notin u^{-1}(r, 1], y \in u^{-1}(r, 1]$ and $x \in v^{-1}(s, 1], y \notin v^{-1}(s, 1]$. Hence (X, T) is T_1 -space.

This completes the proof.

Thus it is seen that $\alpha - T_1(p)$ is a good extension of its topological counter part.
($p = i, ii, iii, iv$)

3.2.11. Theorem : - Let (X, t) be a fuzzy topological space, and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$, then

(1) (X, t) is $\alpha - T_1(i) \Rightarrow (X, I_\alpha(t))$ is T_1 .

(2) (X, t) is $\alpha - T_1(ii) \Rightarrow (X, I_\alpha(t))$ is T_1 .

(3) (X, t) is $\alpha - T_1(iii) \Leftrightarrow (X, I_\alpha(t))$ is T_1 .

Proof : - Let (X, t) be $\alpha - T_1(i)$. We shall prove that $(X, I_\alpha(t))$ is T_1 . Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_1(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 1, u(y) \leq \alpha$ and $v(x) \leq \alpha, v(y) = 1$. Since $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$, and it is clear that $x \in u^{-1}(\alpha, 1], y \notin u^{-1}(\alpha, 1]$ and $x \notin v^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$. Hence $(X, I_\alpha(t))$ is T_1 -space.

Again, suppose that (X, t) is $\alpha - T_1(ii)$ space. We shall prove that $(X, I_\alpha(t))$ is T_1 -space. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_1(ii)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 0, u(y) > \alpha$ and $v(x) > \alpha, v(y) = 0$. Since $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$ and it is clear that $x \notin u^{-1}(\alpha, 1], y \in u^{-1}(\alpha, 1]$ and $x \in v^{-1}(\alpha, 1], y \notin v^{-1}(\alpha, 1]$. Hence $(X, I_\alpha(t))$ is T_1 -Space.

Further, suppose that (X, t) is $\alpha - T_1(iii)$. We shall prove that $(X, I_\alpha(t))$ is T_1 .

Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_1(iii)$, for $\alpha \in I_1, \exists u, v \in t$ such that

$0 \leq u(x) \leq \alpha < u(y) \leq 1$ and $0 \leq v(y) \leq \alpha < v(x) \leq 1$. Since $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$ and it is clear that $x \notin u^{-1}(\alpha, 1], y \in u^{-1}(\alpha, 1]$ and $x \in v^{-1}(\alpha, 1], y \notin v^{-1}(\alpha, 1]$. Hence $(X, I_\alpha(t))$ is T_1 -Space.

Conversely, suppose that $(X, I_\alpha(t))$ is T_1 -space. We shall prove that (X, t) is $\alpha\text{-}\mathcal{T}_1$ (iii). Let $x, y \in X$ with $x \neq y$. Since $(X, I_\alpha(t))$ is T_1 -space so $\exists M, N \in I_\alpha(t)$ such that $x \in M, y \notin M$ and $x \notin N, y \in N$, where $M = u^{-1}(\alpha, 1], N = v^{-1}(\alpha, 1]$, where $u, v \in t$. So it is clear that $u(x) > \alpha, u(y) \leq \alpha$ and $v(x) \leq \alpha, v(y) > \alpha$. This implies that $0 \leq u(y) \leq \alpha < u(x) \leq 1$ and $0 \leq v(x) \leq \alpha < v(y) \leq 1$. Hence (X, t) is $\alpha\text{-}\mathcal{T}_1$ (iii).

Now, we give an example.

3.2.12. Example :- Let $X = \{x, y\}$ and $u, v, w \in I^X$, where u, v, w are defined by $u(x) = 1, u(y) = 0, v(x) = 0.42, v(y) = 0.95, w(x) = 0.15, w(y) = 0.32$. Consider the fuzzy topology t on X generated by $\{0, u, v, w, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.61$, it is clear that (X, t) is not $\alpha\text{-}\mathcal{T}_1$ (i) and (X, t) is not $\alpha\text{-}\mathcal{T}_1$ (ii). Now $I_\alpha(t) = \{X, \phi, \{x\}, \{y\}\}$. Then we see that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is T_1 space.

This completes the proof.

3.2.13. Theorem :- Let (X, t) be a fuzzy topological space, $A \subseteq X$,

$t_A = \{u/A : u \in t\}$, then

- (a) (X, t) is $\alpha\text{-}\mathcal{T}_1$ (i) $\Rightarrow (A, t_A)$ is $\alpha\text{-}\mathcal{T}_1$ (i).
- (b) (X, t) is $\alpha\text{-}\mathcal{T}_1$ (ii) $\Rightarrow (A, t_A)$ is $\alpha\text{-}\mathcal{T}_1$ (ii).
- (c) (X, t) is $\alpha\text{-}\mathcal{T}_1$ (iii) $\Rightarrow (A, t_A)$ is $\alpha\text{-}\mathcal{T}_1$ (iii).
- (d) (X, t) is \mathcal{T}_1 (iv) $\Rightarrow (A, t_A)$ is \mathcal{T}_1 (iv).

Proof : - Suppose (X, t) be an fts and (X, t) is $\alpha - T_1(i)$. We shall prove that (A, t_A) is $\alpha - T_1(i)$. Let $x, y \in A$ with $x \neq y$, then $x, y \in X, x \neq y$, as $A \subseteq X$, Since (X, t) is $\alpha - T_1(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 1, u(y) \leq \alpha$ and $v(x) \leq \alpha, v(y) = 1$. For $A \subseteq X$, we find that $u/A, v/A \in t_A$ and $u/A(x) = 1, u/A(y) \leq \alpha$ and $v/A(x) \leq \alpha, v/A(y) = 1$ as $x, y \in A$. Hence it is clear that (A, t_A) is $\alpha - T_1(i)$.

Similarly (b), (c) and (d) can be proved.

This completes the proof.

3.2.14. Theorem : - Given $(X_i, t_i), i \in \Lambda$ be fuzzy topological space and $X = \prod_{i \in \Lambda} X_i$.

Let t be the product fuzzy topology on X , then

- (a) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - T_1(i) \Leftrightarrow (X, t)$ is $\alpha - T_1(i)$.
- (b) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - T_1(ii) \Leftrightarrow (X, t)$ is $\alpha - T_1(ii)$.
- (c) $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - T_1(iii) \Leftrightarrow (X, t)$ is $\alpha - T_1(iii)$.
- (d) $\forall i \in \Lambda, (X_i, t_i)$ is $T_1(iv) \Leftrightarrow (X, t)$ is $T_1(iv)$.

Proof : - Suppose that $\forall i \in \Lambda, (X_i, t_i)$ is $\alpha - T_1(i)$. We shall prove that (X, t) is $\alpha - T_1(i)$.

Let $x, y \in X$ with $x \neq y$, then $x_i \neq y_i$, for some $i \in \Lambda$. Since (X_i, t_i) is $\alpha - T_1(i)$, for $\alpha \in I_1, \exists u_i, v_i \in t_i, i \in \Lambda$, such that $u_i(x_i) = 1, u_i(y_i) \leq \alpha$ and $v_i(x_i) \leq \alpha, v_i(y_i) = 1$. But we have $\pi_i(x) = x_i$ and $\pi_i(y) = y_i$. Then $u_i(\pi_i(x)) = 1, u_i(\pi_i(y)) \leq \alpha$ and $v_i(\pi_i(x)) \leq \alpha, v_i(\pi_i(y)) = 1$. It follows that $\exists u_i \circ \pi_i, v_i \circ \pi_i \in t$ such that $(u_i \circ \pi_i)(x) = 1, (u_i \circ \pi_i)(y) \leq \alpha$ and $(v_i \circ \pi_i)(x) \leq \alpha, (v_i \circ \pi_i)(y) = 1$. Hence it is clear that (X, t) is $\alpha - T_1(i)$.

Conversely, suppose that (X, t) is $\alpha - T_1(i)$. We shall prove that $(X_i, t_i), i \in \Lambda$ is $\alpha - T_1(i)$. Let fore some $i \in \Lambda, a_i$ be a fixed element in X_i , suppose that $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$. So that A_i is a subset of X , and hence (A_i, t_{A_i}) is also

a subspace of (X, t) . Since (X, t) is $\alpha - T_1(i)$, then (A_i, t_{A_i}) is also $\alpha - T_1(i)$. Now we have A_i is a homeomorphic image of X_i . Hence $(X_i, t_i), i \in \Lambda$, is $\alpha - T_1(i)$.

Similarly (b), (c) and (d) can be proved.

3.2.15. Theorem :-

$$\begin{array}{ccc}
 \text{(a) } (X, t) \text{ is } \alpha - T_1(i) \Rightarrow (X, t) \text{ is } \alpha - T_1(\text{iii}) \Rightarrow (X, t) \text{ is } T_1(\text{iv}). & & \\
 \Downarrow & \Downarrow & \Downarrow \\
 (X, t) \text{ is } \alpha - T_0(i) \Rightarrow (X, t) \text{ is } \alpha - T_0(\text{iii}) \Rightarrow (X, t) \text{ is } T_0(\text{iv}). & & \\
 \\
 \text{(b) } (X, t) \text{ is } \alpha - T_1(\text{ii}) \Rightarrow (X, t) \text{ is } \alpha - T_1(\text{iii}) \Rightarrow (X, t) \text{ is } T_1(\text{iv}) & & \\
 \Downarrow & \Downarrow & \Downarrow \\
 (X, t) \text{ is } \alpha - T_0(\text{ii}) \Rightarrow (X, t) \text{ is } \alpha - T_0(\text{iii}) \Rightarrow (X, t) \text{ is } T_0(\text{iv}). & &
 \end{array}$$

Proof :- The proofs of (a) and (b) are easy.

3.2.16. Theorem:- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be a one-one, onto and open map then,

- (a) (X, t) is $\alpha - T_1(i) \Rightarrow (Y, s)$ is $\alpha - T_1(i)$.
- (b) (X, t) is $\alpha - T_1(\text{ii}) \Rightarrow (Y, s)$ is $\alpha - T_1(\text{ii})$.
- (c) (X, t) is $\alpha - T_1(\text{iii}) \Rightarrow (Y, s)$ is $\alpha - T_1(\text{iii})$.
- (d) (X, t) is $T_1(\text{iv}) \Rightarrow (Y, s)$ is $T_1(\text{iv})$.

Proof :- Suppose (X, t) be $\alpha - T_1(i)$. We shall prove that (Y, s) is $\alpha - T_1(i)$. Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto then $\exists x_1, x_2 \in X$ with $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is

one-one . Again since (X , t) is $\alpha - T_1(i)$, for $\alpha \in I_1$, $\exists u , v \in t$ such that $u(x_1) = 1$, $u(x_2) \leq \alpha$ and $v(x_1) \leq \alpha$, $v(x_2) = 1$.

$$\begin{aligned} \text{Now } f(u) (y_1) &= \{ \text{Sup } u(x_1) \quad ; f(x_1) = y_1 \} \\ &= 1. \end{aligned}$$

$$\begin{aligned} f(u) (y_2) &= \{ \text{Sup } u(x_2) \quad ; f(x_2) = y_2 \} \\ &\leq \alpha \end{aligned}$$

$$\begin{aligned} \text{and } f(v) (y_1) &= \{ \text{Sup } v(x_1) \quad ; f(x_1) = y_1 \} \\ &\leq \alpha \end{aligned}$$

$$\begin{aligned} f(v) (y_2) &= \{ \text{Sup } v(x_2) \quad ; f(x_2) = y_2 \} \\ &= 1 \end{aligned}$$

Since f is open then $f(u), f(v) \in s$. Now it is clear that $\exists f(u) , f(v) \in s$ such that $f(u) (y_1) = 1$, $f(u) (y_2) \leq \alpha$ and $f(v) (y_1) \leq \alpha$, $f(v) (y_2) = 1$. Hence (Y , s) is $\alpha - T_1(i)$.

Similarly (b) , (c) and (d) can be proved .

3.2.17. Theorem:- Let (X , t) and (Y , s) be two fuzzy topological spaces and

$f : X \longrightarrow Y$ be a continuous and one-one map then ,

$$(a) (Y , s) \text{ is } \alpha - T_1(i) \Rightarrow (X , t) \text{ is } \alpha - T_1(i) .$$

$$(b) (Y , s) \text{ is } \alpha - T_1(ii) \Rightarrow (X , t) \text{ is } \alpha - T_1(ii) .$$

$$(c) (Y , s) \text{ is } \alpha - T_1(iii) \Rightarrow (X , t) \text{ is } \alpha - T_1(iii) .$$

$$(d) (Y , s) \text{ is } T_1(iv) \Rightarrow (X , t) \text{ is } T_1(iv) .$$

Proof :- Suppose (Y , s) be $\alpha - T_1(i)$. We shall prove that (X , t) is $\alpha - T_1(i)$. Let $x_1, x_2 \in X$ with $x_1 \neq x_2$, then $f(x_1) \neq f(x_2)$ in Y as f is one-one . Since (Y , s) is $\alpha - T_1(i)$, for $\alpha \in I_1$, $\exists u , v \in t$ such that $u(f(x_1)) = 1$, $u(f(x_2)) \leq \alpha$ and $v(f(x_1)) \leq \alpha$, $v(f(x_2)) = 1$. This implies that $f^{-1}(u) (x_1) = 1$, $f^{-1}(u) (x_2) \leq \alpha$ and $f^{-1}(v) (x_1) \leq \alpha$, $f^{-1}(v) (x_2) = 1$, since f is

continuous and $u, v \in s$ then $f^{-1}(u), f^{-1}(v) \in t$. Now it is clear that $\exists f^{-1}(u), f^{-1}(v) \in t$ such that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) \leq \alpha$ and $f^{-1}(v)(x_1) \leq \alpha, f^{-1}(v)(x_2) = 1$. Hence (X, t) is $\alpha - T_1(i)$.

Similarly (b), (c) and (d) can be proved.

Chapter : -4

T_2 Topological Space

4. Introduction:-

In this chapter, we introduce and study some T_2 properties in fuzzy topological spaces and obtain their several features.

4.1. Definition:-


Let (X, t) be a fuzzy topological space and $\alpha \in I_1$.

(a) (X, t) is $\alpha - T_2(i)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) = 1 = v(y)$ and $u \cap v \leq \alpha$.

(b) (X, t) is $\alpha - T_2(ii)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$.

(c) (X, t) is $\alpha - T_2(iii)$ space $\Leftrightarrow \forall x, y \in X$ with $x \neq y, \exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$.

4.2. Lemma :- The following implication are true .



Proof:- Let (X, t) be a fuzzy topological space and (X, t) is $\alpha - T_2(i)$. We shall prove that (X, t) is $\alpha - T_2(iii)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_2(i)$, for $\alpha \in I_1$,

$\exists u, v \in \tau$ such that $u(x) = 1 = v(y)$ and $u \cap v \leq \alpha$. So it is clear that $u(x) > \alpha$, $v(y) > \alpha$ and $u \cap v \leq \alpha$. Hence it is clear that (X, τ) is $\alpha - T_2(\text{iii})$.

Next, suppose that (X, τ) is $\alpha - T_2(\text{ii})$. We shall prove that (X, τ) is $\alpha - T_2(\text{iii})$. Let $x, y \in X$ with $x \neq y$. Since (X, τ) is $\alpha - T_2(\text{ii})$, for $\alpha \in I_1$, $\exists u, v \in \tau$ such that $u(x) > \alpha$, $v(y) > \alpha$ and $u \cap v = 0$. Now it is clear that $u(x) > \alpha$, $v(y) > \alpha$ and $u \cap v \leq \alpha$. Hence (X, τ) is $\alpha - T_2(\text{iii})$.

Now, we give some examples to show the non implication among $\alpha - T_2(\text{i})$, $\alpha - T_2(\text{ii})$ and $\alpha - T_2(\text{iii})$.

4.3. Example :- Let $X = \{x, y\}$, and $u, v \in I^X$, where u and v are defined by $u(x) = 0.7$, $u(y) = 0$, $v(x) = 0$ and $v(y) = 0.7$. Consider the fuzzy topology τ on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$ it is clear that (X, τ) is $\alpha - T_2(\text{ii})$ but (X, τ) is not $\alpha - T_2(\text{i})$.

4.4. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1 = v(y)$, and $u(y) = 0.4 = v(x)$. Consider the fuzzy topology τ on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$, we see that (X, τ) is $\alpha - T_2(\text{i})$ but (X, τ) is not $\alpha - T_2(\text{ii})$.

4.5. Examples :- Let $X = \{x, y\}$, and $u, v \in I^X$, where u, v are defined by $u(x) = 0.9$, $u(y) = 0.3$, $v(x) = 0.2$ and $v(y) = 0.9$. Consider the fuzzy topology τ on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$, we get (X, τ) is $\alpha - T_2(\text{iii})$, but (X, τ) is not $\alpha - T_2(\text{i})$ and (X, τ) is not $\alpha - T_2(\text{ii})$.

4.6. Theorem :- If $0 \leq \alpha \leq \beta < 1$, then

(a) (X, t) is $\alpha - T_2(i) \Rightarrow (X, t)$ is $\beta - T_2(i)$.

(b) (X, t) is $\beta - T_2(ii) \Rightarrow (X, t)$ is $\alpha - T_2(ii)$.

(c) (X, t) is $0 - T_2(ii) \Leftrightarrow (X, t)$ is $0 - T_2(iii)$.

Proof :- Suppose (X, t) is $\alpha - T_2(i)$. We shall prove that (X, t) is $\beta - T_2(i)$. Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_2(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 1 = v(y)$ and $u \cap v \leq \alpha$. This implies that $u(x) = 1 = v(y)$ and $u \cap v \leq \beta$ as $0 \leq \alpha \leq \beta < 1$. Hence (X, t) is $\beta - T_2(i)$.

4.7. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1, u(y) = 0.6, v(x) = 0.7, v(y) = 1$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.3$ and $\beta = .08$, we get (X, t) is $\beta - T_2(i)$, but (X, t) is not $\alpha - T_2(i)$.

Next, suppose that (X, t) is $\beta - T_2(ii)$. We shall prove that (X, t) is $\alpha - T_2(ii)$. Let $x, y \in X$, with $x \neq y$. Since (X, t) is $\beta - T_2(ii)$, for $\beta \in I_1, \exists u, v \in t$ such that $u(x) > \beta, v(y) > \beta$ and $u \cap v = 0$. This implies that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$. as $0 \leq \alpha \leq \beta < 1$. Hence it is clear that (X, t) is $\alpha - T_2(ii)$.

4.8. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.6, u(y) = 0, v(x) = 0$ and $v(y) = 0.7$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$ and $\beta = 0.9$, we get (X, t) is $\alpha - T_2(ii)$, but (X, t) is not $\beta - T_2(ii)$.

Finally, suppose that (X, t) is $0 - T_2(ii)$. We shall prove that (X, t) is $0 - T_2(iii)$. Let $x, y \in X$, with $x \neq y$. Since (X, t) is $0 - T_2(ii)$, $\exists u, v \in t$ such that $u(x) > 0, v(y) > 0$

and $u \cap v = 0$. This implies that $u(x) > 0, v(y) > 0$ and $u \cap v \leq 0$. Hence it is clear that (X, τ) is $0 - T_2(\text{iii})$.


Conversely, suppose that (X, τ) is $0 - T_2(\text{iii})$. We shall prove that (X, τ) is $0 - T_2(\text{ii})$. Let $x, y \in X$, with $x \neq y$. Since (X, τ) is $0 - T_2(\text{iii})$, $\exists u, v \in \tau$ such that $u(x) > 0, v(y) > 0$ and $u \cap v \leq 0$. This implies that $u(x) > 0, v(y) > 0$ and $u \cap v = 0$. Hence it is clear that (X, τ) is $0 - T_2(\text{ii})$.

This completes the proof.

4.9. Theorem :- Let (X, T) be a topological space. Consider the following statements:

- (1) (X, T) is a T_2 - space .
- (2) $(X, \omega(T))$ is $\alpha - T_2(\text{i})$ space .
- (3) $(X, \omega(T))$ is $\alpha - T_2(\text{ii})$ space .
- (5) $(X, \omega(T))$ is $\alpha - T_2(\text{iii})$ space.

Then the following implications are true.



Proof :- Let (X, T) be a T_2 - space. We shall prove that $(X, \omega(T))$ is $\alpha - T_2(\text{i})$.

Let $x, y \in X$, with $x \neq y$. Since (X, T) is T_2 - space, $\exists U, V \in T$ such that $x \in U, y \in V$ and $U \cap V = \phi$. From the definition of lower semi continuous function $1_U, 1_V \in \omega(T)$ and $1_U(x) = 1, 1_V(y) = 1$ and $1_U \cap 1_V = 0$. If $1_U \cap 1_V \neq 0$, then $\exists z \in X$ such that $(1_U \cap 1_V)(z) \neq 0 \Rightarrow 1_U(z) \neq 0, 1_V(z) \neq 0, \Rightarrow z \in U, z \in V \Rightarrow z \in U \cap V \Rightarrow U \cap V \neq \phi$, a contradiction.

So that $1_U \cap 1_V = 0$, and consequently $(X, \omega(T))$ is $\alpha - T_2(\text{i})$. Also, we see that $(X, \omega(T))$ is $\alpha - T_2(\text{ii})$.

Further, it is easy to show that (2) \Rightarrow (4) and (3) \Rightarrow (4) .

We, therefore prove that (4) \Rightarrow (1) .

Suppose that $(X, \omega(T))$ is $\alpha - T_2$ (iii) space. We shall prove that (X, T) is T_2 - space. Let $x, y \in X$ with $x \neq y$. Since $(X, \omega(T))$ is $\alpha - T_2$ (iii) , for some $\alpha \in I_1$, then $\exists u, v \in \omega(T)$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. Now we have that $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in T$, $\alpha \in I_1$ and $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$. Moreover $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$. For if $z \in u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1]$, then $z \in u^{-1}(\alpha, 1]$ and $z \in v^{-1}(\alpha, 1] \Rightarrow u(z) > \alpha$ and $v(z) > \alpha \Rightarrow (u \cap v)(z) > \alpha$, a contradiction as $(u \cap v)(z) \leq \alpha$. Hence (X, T) is T_2 - space. This completes the proof.

Thus it is seen that $\alpha - T_2(p)$ is a good extension of its topological counter part
($p = i, ii, iii,$)

4.10. Theorem :- Let (X, t) be a fuzzy topological space, and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$. Then

(a) (X, t) is $\alpha - T_2(i) \Rightarrow (X, I_\alpha(t))$ is T_2 .

(b) (X, t) is $\alpha - T_2(ii) \Rightarrow (X, I_\alpha(t))$ is T_2 .

(c) (X, t) is $\alpha - T_2(iii) \Leftrightarrow (X, I_\alpha(t))$ is T_2 .

Proof :- Consider (X, t) be a fts and (X, t) is $\alpha - T_2(i)$. We shall prove that $(X, I_\alpha(t))$ is T_2 . Let $x, y \in X$ with $x \neq y$. Since (X, t) is $\alpha - T_2(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) = 1, v(y) = 1$ and $u \cap v \leq \alpha$. But for every $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$ and also $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$, as $u \cap v \leq \alpha$. Hence it is clear that $(X, I_\alpha(t))$ is T_2 - space .

Next, suppose that (X, t) is α - $T_2(ii)$. We shall prove that $(X, I_\alpha(t))$ is T_2 -space.

Let $x, y \in X$, with $x \neq y$. Since (X, t) is α - $T_2(ii)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v = 0$. But for every $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$. So we have $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$, as $u \cap v = 0$. Hence it is clear that $(X, I_\alpha(t))$ is T_2 -space.

Further, suppose that (X, t) is α - $T_2(iii)$. We shall prove that $(X, I_\alpha(t))$ is T_2 -space.

Let $x, y \in X$ with $x \neq y$. Since (X, t) is α - $T_2(iii)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. But for every $\alpha \in I_1, u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in I_\alpha(t)$. So we have $x \in u^{-1}(\alpha, 1], y \in v^{-1}(\alpha, 1]$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$, as $u \cap v \leq \alpha$. Hence it is clear that $(X, I_\alpha(t))$ is T_2 -space.

Conversely, suppose that $(X, I_\alpha(t))$ is T_2 -space. We shall prove that (X, t) is α - $T_2(iii)$. Let $x, y \in X$ with $x \neq y$. Since $(X, I_\alpha(t))$ is T_2 -space, $\exists U, V \in I_\alpha(t)$, such that $x \in U, y \in V$ and $U \cap V = \phi$. Again since $U, V \in I_\alpha(t)$, so we get $u, v \in t$ such that $U = u^{-1}(\alpha, 1], V = v^{-1}(\alpha, 1]$. This implies that $u(x) > \alpha, v(y) > \alpha$ and $u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi \Rightarrow (u \cap v)^{-1}(\alpha, 1] = \phi$ ie $u \cap v \leq \alpha$. So, we see that (X, t) is α - $T_2(iii)$.

Now we give an example.

4.11. Example :- Let $X = \{x, y\}$, and $u, v \in I^X$, where u, v are defined by $u(x) = 0.8, u(y) = 0.2, v(x) = 0.3$ and $v(y) = 0.7$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.5$, we see that (X, t) is not α - $T_2(i)$ and (X, t) is not α - $T_2(ii)$. Now $I_\alpha(t) = \{X, \phi, \{x\}, \{y\}\}$. Then we see that $I_\alpha(t)$ is a topology on X and $(X, I_\alpha(t))$ is T_2 -space.

This completes the proof.

4.12. Theorem :- Let (X, t) be a fuzzy topological space and $A \subseteq X$,

$t_A = \{ u/A : u \in t \}$, then

(a) (X, t) is an $\alpha - T_2(i) \Rightarrow (A, t_A)$ is an $\alpha - T_2(i)$.

(b) (X, t) is an $\alpha - T_2(ii) \Rightarrow (A, t_A)$ is an $\alpha - T_2(ii)$.

(c) (X, t) is an $\alpha - T_2(iii) \Rightarrow (A, t_A)$ is an $\alpha - T_2(iii)$.

Proof :- Suppose (X, t) be an $\alpha - T_2(iii)$. We shall prove that (A, t_A) is an $\alpha - T_2(iii)$.

Let $x, y \in A$, with $x \neq y$, then $x, y \in X$, with $x \neq y$. Since (X, t) is $\alpha - T_2(iii)$, for $\alpha \in I_1$, $\exists u, v \in t$ such that $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. But we have $u/A, v/A \in t_A$ for every $u, v \in t$. This implies that $u/A(x) > \alpha, v/A(y) > \alpha$ and $u/A \cap v/A \leq \alpha$. Hence it is clear that (A, t_A) is $\alpha - T_2(iii)$.

Similarly, other cases can be proved.

4.13. Theorem :- Given $\{ (X_i, t_i), i \in \Lambda \}$ be fuzzy topological spaces and $X = \prod_{i \in \Lambda} X_i$ and t be the product fuzzy topology on X . Then

(a) $\forall i \in \Lambda, (X_i, t_i)$ is an $\alpha - T_2(i) \Leftrightarrow (X, t)$ is an $\alpha - T_2(i)$.

(b) $\forall i \in \Lambda, (X_i, t_i)$ is an $\alpha - T_2(ii) \Leftrightarrow (X, t)$ is an $\alpha - T_2(ii)$.

(c) $\forall i \in \Lambda, (X_i, t_i)$ is an $\alpha - T_2(iii) \Leftrightarrow (X, t)$ is an $\alpha - T_2(iii)$.

Proof :- Suppose $\forall i \in \Lambda, (X_i, t_i)$ be an $\alpha - T_2(iii)$. We shall prove that (X, t) is $\alpha - T_2(iii)$.

Let x, y be two distinct points in $X = \prod_{i \in \Lambda} X_i$, then there exist an $x_i \neq y_i$ in X_i . Since (X_i, t_i) is an $\alpha - T_2(iii)$, for $\alpha \in I_1$, $\exists u_i, v_i \in t_i$ such that $u_i(x_i) > \alpha, v_i(y_i) > \alpha$ and $u_i \cap v_i \leq \alpha$.

But we have $\pi_i(x) = x_i, \pi_i(y) = y_i$, then $u_i(\pi_i(x)) > \alpha, v_i(\pi_i(y)) > \alpha$ and $(u_i \cap v_i) \circ \pi_i \leq \alpha$.



Hence $(u_i \circ \pi_i)(x) > \alpha, (v_i \circ \pi_i)(y) > \alpha$ and $(u_i \circ \pi_i) \cap (v_i \circ \pi_i) \leq \alpha$. Put $u = u_i \circ \pi_i$,

$v = v_i \circ \pi_i$, then $u, v \in \mathcal{t}$ with $u(x) > \alpha, v(y) > \alpha$ and $u \cap v \leq \alpha$. Hence it is clear that (X, \mathcal{t}) is $\alpha - T_2(\text{iii})$.

Conversely, suppose that (X, \mathcal{t}) is $\alpha - T_2(\text{iii})$. We shall prove that (X_i, \mathcal{t}_i) is $\alpha - T_2(\text{iii})$, for $i \in \Lambda$. For some $i \in \Lambda$, let a_i be a fixed element in X_i . Suppose that $A_i = \{x \in X = \prod_{i \in \Lambda} X_i : x_j = a_j \text{ for some } i \neq j\}$. Then A_i is a subsets of X and therefore (A_i, \mathcal{t}_{A_i}) is a subspace of (X, \mathcal{t}) . Since (X, \mathcal{t}) is $\alpha - T_2(\text{iii})$ space. Then, we have also (A_i, \mathcal{t}_{A_i}) is also $\alpha - T_2(\text{iii})$ space. Further more, A_i is homeomorphic image of X_i . Hence it is clear that (X_i, \mathcal{t}_i) is $\alpha - T_2(\text{iii})$ space.

The proofs for (a) and (b) are similar.

4.14. Theorem:-



Proof :- The proofs of (a) and (b) are easy.

4.15. Theorem:- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be a one-one, onto and open map, then

$$(a) (X, t) \text{ is } \alpha\text{-}T_2(i) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_2(i).$$

$$(b) (X, t) \text{ is } \alpha\text{-}T_2(ii) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_2(ii).$$

$$(c) (X, t) \text{ is } \alpha\text{-}T_2(iii) \Rightarrow (Y, s) \text{ is } \alpha\text{-}T_2(iii).$$

Proof:- Suppose (X, t) is $\alpha\text{-}T_2(i)$. We shall prove that (Y, s) is $\alpha\text{-}T_2(i)$. Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto then, $\exists x_1, x_2 \in X$ with $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one. Again since (X, t) is $\alpha\text{-}T_2(i)$, for $\alpha \in I_1, \exists u, v \in t$ such that $u(x_1) = 1 = v(x_2)$ and $u \cap v \leq \alpha$.

$$\begin{aligned} \text{Now } f(u)(y_1) &= \{ \text{Sup } u(x_1) : f(x_1) = y_1 \} \\ &= 1 \end{aligned}$$

$$f(v)(y_2) = \{ \text{Sup } v(x_2) : f(x_2) = y_2 \}$$

$$\text{and } f(u \cap v)(y_1) = \{ \text{Sup } (u \cap v)(x_1) : f(x_1) = y_1 \}$$

$$f(u \cap v)(y_2) = \{ \text{Sup } (u \cap v)(x_2) : f(x_2) = y_2 \}$$

$$\text{Hence } f(u \cap v) \leq \alpha \Rightarrow f(u) \cap f(v) \leq \alpha.$$

Since f is open then $f(u), f(v) \in s$. Now it is clear that $\exists f(u), f(v) \in s$ such that $f(u)(y_1) = 1, f(v)(y_2) = 1$ and $f(u) \cap f(v) \leq \alpha$. Hence (Y, s) is $\alpha\text{-}T_2(i)$.

Similarly (b) and (c) can be proved.

4.16. Theorem :- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be continuous and one-one map then,

$$(a) (Y, s) \text{ is } \alpha\text{-}T_2(i) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_2(i).$$

$$(b) (Y, s) \text{ is } \alpha\text{-}T_2(ii) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_2(ii).$$

$$(c) (Y, s) \text{ is } \alpha\text{-}T_2(iii) \Rightarrow (X, t) \text{ is } \alpha\text{-}T_2(iii).$$

Proof:- Suppose (Y, s) is $\alpha - T_2(i)$. We shall prove that (X, t) is $\alpha - T_2(i)$. Let $x_1, x_2 \in X$ with $x_1 \neq x_2$, then $f(x_1) \neq f(x_2)$ in Y , as f is one-one. Since (Y, s) is $\alpha - T_2(i)$, for $\alpha \in I_1$, then $\exists u, v \in s$ such that $u(f(x_1)) = 1 = v(f(x_2))$ and $u \cap v \leq \alpha$. This implies that $f^{-1}(u)(x_1) = 1$, $f^{-1}(v)(x_2) = 1$ and $f^{-1}(u \cap v) \leq \alpha$ ie $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$, since $u, v \in s$ and f is continuous then $f^{-1}(u), f^{-1}(v) \in t$. Now it is clear that $\exists f^{-1}(u), f^{-1}(v) \in t$ such that $f^{-1}(u)(x_1) = 1$, $f^{-1}(v)(x_2) = 1$ and $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$. Hence (X, t) is $\alpha - T_2(i)$.

Similarly (b) and (c) can be proved.

Chapter : -5

Regular Fuzzy Topological Space

5. Introduction:-

In this chapter, we introduce and study some Regular property in fuzzy topological spaces and obtain their several features.

5.1. Definition:- Let (X, τ) be a fuzzy topological space and $\alpha \in I_1$.


(a) (X, τ) is an α -FR(i) space $\Leftrightarrow \forall w \in \tau^c, \forall x \in X$, with $w(x) < 1, \exists u, v \in \tau$ such that $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$.

(b) (X, τ) is an α -FR(ii) space $\Leftrightarrow \forall w \in \tau^c, \forall x \in X$, with $w(x) < 1, \exists u, v \in \tau$ such that $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$.

(c) (X, τ) is an α -FR(iii) space $\Leftrightarrow \forall w \in \tau^c, \forall x \in X$, with $w(x) = 0, \exists u, v \in \tau$ such that $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$.

(d) (X, τ) is an α -FR(iv) space $\Leftrightarrow \forall w \in \tau^c, \forall x \in X$, with $w(x) = 0, \exists u, v \in \tau$ such that $u(x) > \alpha, v(y) = 1, y \in w^{-1}\{1\}, u \cap v \leq \alpha$.

5.2. Theorem:- The following implications are true :



Proof:- First , suppose that (X , t) is α - FR(i). We shall prove that (X , t) is α - FR(ii) .

Let $w \in t^c$, $x \in X$, with $w(x) < 1$. Since (X , t) is α -FR(i) , for $\alpha \in I_1$, $\exists u , v \in t$ such that $u(x) = 1$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Now, we see that $u(x) > \alpha$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Hence it is clear that (X , t) is α -FR(ii) .

Next, suppose that (X , t) is α -FR(i) . We shall prove that (X , t) is α - FR(iii) .

Let $w \in t^c$, $x \in X$, with $w(x) = 0$. Then we have $w(x) < 1$. Since (X , t) is α -FR(i) , for $\alpha \in I_1$ $\exists u , v \in t$ such that $u(x) = 1$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Now it is clear that (X , t) is α -FR(iii).

Again, suppose that (X , t) is α -FR(ii) . We shall prove that (X , t) is α -FR(iv).

Let $w \in t^c$, $x \in X$, with $w(x) = 0$, Then clearly $w(x) < 1$. Since (X , t) is α -FR(ii) , for $\alpha \in I_1$, $\exists u , v \in t$ such that $u(x) > \alpha$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Hence it is clear that (X , t) is α -FR(iv) .

Further, suppose that (X , t) is α - FR(iii) . We shall prove that (X , t) is α -FR(iv).

Let $w \in t^c$ and $x \in X$, with $w(x) = 0$. Since (X , t) is α - Fr(iii) , for $\alpha \in I_1$, $\exists u , v \in t$ such that $u(x) = 1$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Hence one can see that (X , t) is α -FR(iv).

Now , we give some examples to show the non implication among α - FR(i) , α - FR(ii) , α - FR(iii) and α - FR(iv) .

5.3. Example :- Let $X = \{ x , y \}$ and $u , v \in I^X$, where u , v are defined by $u(x) = 0.9$, $u(y) = 0$, $v(x) = 0.5$ and $v(y) = 1$. Consider the fuzzy topology t on X generated by

$\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $w = 1 - u$ and $\alpha = 0.7$, we see that (X, t) is α -FR(ii) but (X, t) is not α -FR(i).

5.4. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 0.2$, $u(y) = 0.3$, $v(x) = 0.3$, $v(y) = 0.2$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $w = 1 - u$ and $\alpha = 0.5$, we see that (X, t) is α -FR(iii) and (X, t) is α -FR(iv), but (X, t) is not α -FR(ii). As there do not exist any $u, v \in t$ such that $u(x) > \alpha$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$.

5.5. Example :- Let $X = \{x, y\}$ and $u, v, w \in I^X$, where u, v and w are defined by $u(x) = 0.9$, $u(y) = 0$, $v(x) = 0.5$, $v(y) = 1$, $w(x) = 1$, $w(y) = 0$. Consider the fuzzy topology t on X generated by $\{0, u, v, w, 1\} \cup \{\text{Constants}\}$. For $\alpha = 0.6$ and $p = 1 - w$, it is seen that (X, t) is α -FR(iv) but (X, t) is not α -FR(iii).

This completes the proof.

5.6. Theorem :- Let $0 \leq \alpha \leq \beta < 1$, then

- (a) (X, t) is α -FR(i) $\Rightarrow (X, t)$ is β -FR(i).
- (b) (X, t) is α -FR(iii) $\Rightarrow (X, t)$ is β -FR(iii).

Proof :- First, suppose that (X, t) is α -FR(i). We shall prove that (X, t) is β -FR(i). Let $w \in t^c$ and $x \in X$ with $w(x) < 1$. Since (X, t) is α -FR(i), for $\alpha \in I_1$, $\exists u, v \in t$ such that $u(x) = 1$, $v(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Since $\alpha \leq \beta$, then $u \cap v \leq \beta$. So it is observed that (X, t) is β -FR(i).

Next, suppose that (X, t) is α -FR(iii). We shall prove that (X, t) is β -FR(iii). Let $w \in t^c$, $x \in X$ with $w(x) = 0$. Since (X, t) is α -FR(iii), for $\alpha \in I_1$, $\exists u, v \in t$ such that $u(x) = 1$, $u(y) = 1$, $y \in w^{-1}\{1\}$ and $u \cap v \leq \alpha$. Since $0 \leq \alpha \leq \beta < 1$, then $u \cap v \leq \beta$. Now

it can be written as $\forall w \in t^c, \forall x \in X$, with $w(x) = 0, \exists u, v \in t$ such that $u(x) = 1, v(y) = 1, y \in w^{-1}\{1\}$ and $u \cap v \leq \beta$. Hence it is clear that (X, t) is β -FR(iii).

Now, we give an example.

5.7. Example :- Let $X = \{x, y\}$ and $u, v \in I^X$, where u, v are defined by $u(x) = 1, u(y) = 0, v(x) = 0.7, v(y) = 1$. Consider the fuzzy topology t on X generated by $\{0, u, v, 1\} \cup \{\text{Constants}\}$. For $w = 1 - u, \alpha = 0.75, \beta = 0.6$. We see that (X, t) is β -FR(i) and (X, t) is β -FR(iii), but (X, t) is not α -FR(i) and (X, t) is not α -FR(iii).

5.8. Theorem :- Let (X, t) be a fuzzy topological space and

$$I_\alpha(t) = \{u^{-1}(\alpha, 1] : u \in t\}, \text{ then}$$

$$(X, t) \text{ is } 0\text{-FR(i)} \Rightarrow (X, I_0(t)) \text{ is Regular.}$$

Proof :- Consider (X, t) be a 0 -FR(i). We shall prove that $(X, I_0(t))$ is Regular. Let V be a closed set in $I_0(t)$ and $x \in X$ such that $x \notin V$. Then $V^c \in I_0(t)$ and $x \in V^c$. So by the definition of $I_0(t)$, there exists an $u \in t$ such that $V^c = u^{-1}(0, 1]$ ie $u(x) > 0$. Since $u \in t$, then u^c is closed fuzzy set in t and $u^c(x) < 1$. Since (X, t) is 0 -FR(i), then $\exists v, w \in t$ such that $v(x) = 1, w \geq 1_{(u^c)^{-1}\{1\}}, v \cap w = 0$.

(a) Since $v, w \in t$ then $v^{-1}(0, 1], w^{-1}(0, 1] \in I_0(t)$ and $x \in v^{-1}(0, 1]$

(b) Since $w \geq 1_{(u^c)^{-1}\{1\}}$ then $w^{-1}(0, 1] \supseteq (1_{(u^c)^{-1}\{1\}})^{-1}(0, 1]$.

(c) And $v \cap w = 0$, mean $(v \cap w)^{-1}(0, 1] = v^{-1}(0, 1] \cap w^{-1}(0, 1] = \phi$.

Now, we have

$$\begin{aligned}
(1_{(u^c)^{-1}\{1}})^{-1}(0, 1] &= \{x : 1_{(u^c)^{-1}\{1}}(x) \in (0, 1] \} \\
&= \{x : 1_{(u^c)^{-1}\{1}}(x) = 1 \} \\
&= \{x : x \in (u^c)^{-1}\{1\} \} \\
&= \{x : u^c(x) = 1 \} \\
&= \{x : u(x) = 0 \} \\
&= \{x : x \notin V^c \} \\
&= \{x : x \in V \} \\
&= V
\end{aligned}$$

Put $W = v^{-1}(0, 1]$ and $W^* = w^{-1}(0, 1]$, then $x \in W$, $W^* \supseteq V$ and $W \cap W^* = \phi$.

Hence it is clear that $(X, I_0(t))$ is Regular.

5.9. Theorem :- Let (X, t) be a fuzzy topological space $A \subseteq X$, and $t_A = \{u/A : u \in t\}$,

then

$$1_{((u/A)^c)^{-1}\{1}}(x) = (1_{(u^c)^{-1}\{1}}/A)(x)$$

Proof :- Let w be a closed fuzzy set in t_A ie $w \in t_A^c$, then $u/A = w^c$, where $u \in t$.

Now we have

$$\begin{aligned}
1_{((u/A)^c)^{-1}\{1}}(x) &= \begin{cases} 0 & \text{if } x \notin ((u/A)^c)^{-1}\{1\} \\ 1 & \text{if } x \in ((u/A)^c)^{-1}\{1\} \end{cases} \\
&= \begin{cases} 0 & \text{if } x \notin \{y : (u/A)^c(y) = 1\} \\ 1 & \text{if } x \in \{y : (u/A)^c(y) = 1\} \end{cases} \\
&= \begin{cases} 0 & \text{if } (u/A)^c(x) < 1 \\ 1 & \text{if } (u/A)^c(x) = 1 \end{cases} \\
&= \begin{cases} 0 & \text{if } w(x) < 1 \\ 1 & \text{if } w(x) = 1 \end{cases}
\end{aligned}$$

$$\begin{aligned}
\text{Again } 1_{(u^c)^{-1}\{1\}}(x) &= \begin{cases} 0 & \text{if } x \notin (u^c)^{-1}\{1\} \\ 1 & \text{if } x \in (u^c)^{-1}\{1\} \end{cases} \\
&= \begin{cases} 0 & \text{if } x \notin \{y : u^c(y) = 1\} \\ 1 & \text{if } x \in \{y : u^c(y) = 1\} \end{cases} \\
&= \begin{cases} 0 & \text{if } u^c(x) < 1 \\ 1 & \text{if } u^c(x) = 1 \end{cases}
\end{aligned}$$

$$\begin{aligned}
\text{Now } (1_{(u^c)^{-1}\{1\}}/A)(x) &= \begin{cases} 0 & \text{if } (u^c/A)(x) < 1 \\ 1 & \text{if } (u^c/A)(x) = 1 \end{cases} \\
&= \begin{cases} 0 & \text{if } (u/A)^c(x) < 1 \\ 1 & \text{if } (u/A)^c(x) = 1 \end{cases} \\
&= \begin{cases} 0 & \text{if } w(x) < 1 \\ 1 & \text{if } w(x) = 1 \end{cases}
\end{aligned}$$

Hence it is clear that $1_{((u/A)^c)^{-1}\{1\}}(x) = (1_{(u^c)^{-1}\{1\}}/A)(x)$.

5.10. Theorem :- Let (X, t) be a fuzzy topological space and $A \subseteq X$ and

$t_A = \{u/A : u \in t\}$, then

- (a) (X, t) is α -FR(i) $\Rightarrow (A, t_A)$ is α -FR(i).
- (b) (X, t) is α -FR(ii) $\Rightarrow (A, t_A)$ is α -FR(ii).
- (c) (X, t) is α -FR(iii) $\Rightarrow (A, t_A)$ is α -FR(iii).
- (d) (X, t) is α -FR(iv) $\Rightarrow (A, t_A)$ is α -FR(iv).

Proof :- Let (X, t) be α -FR(i). We shall prove that (A, t_A) is α -FR(i). Let w be a closed fuzzy set in t_A , and $x^* \in A$ such that $w(x^*) < 1$. This implies that $w^c \in t_A$ and $w^c(x^*) > 0$. So there exists an $u \in t$ such that $u/A = w^c$ and clearly u^c is closed in t and $u^c(x^*) = (u/A)^c(x^*) = w(x^*) < 1$, ie $u^c(x^*) < 1$. Since (X, t) is α -FR(i), for $\alpha \in I_1, \exists v, v^* \in t$

such that $v(x^*) = 1, v^* \geq 1_{(u^c)^{-1}\{1\}}$ and $v \cap v^* \leq \alpha$. Since $v, v^* \in t$, then $v/A, v^*/A \in t_A$ and $v/A(x^*) = 1, v^*/A \geq (1_{(u^c)^{-1}\{1\}}/A)$ and $v/A \cap v^*/A = (v \cap v^*)/A \leq \alpha$.


But $1_{(u^c)^{-1}\{1\}}/A = 1_{((u/A)^c)^{-1}\{1\}} = 1_{w^{-1}\{1\}}$, then $v^*/A \geq 1_{w^{-1}\{1\}}$. Hence it is clear that (A, t_A) is α -FR(i).

The proofs of (b), (c) and (d) are similar.

5.11. Theorem :- Let (X, T) be a topological space. Consider the following statements .

- (1) (X, T) is a Regular space.
- (2) $(X, \omega(T))$ is α -FR(i) .
- (3) $(X, \omega(T))$ is α -FR(ii) .
- (4) $(X, \omega(T))$ is α -FR(iii) .
- (5) $(X, \omega(T))$ is α -FR(iv) .

Then



Proof :- First , suppose that (X, T) be regular space . We shall prove that $(X, \omega(T))$ is α -FR(i) . Let w be a closed fuzzy set in $\omega(T)$ and $x \in X$ such that $w(x) < 1$, then $w^c \in \omega(T)$ and $w^c(x) > 0$. Now we have $(w^c)^{-1}(0, 1] \in T, x \in (w^c)^{-1}(0, 1]$. Also it is clear that $[(w^c)^{-1}(0, 1]]^c = w^{-1}\{1\}$ be a closed in T and $x \notin w^{-1}\{1\}$. Since (X, T) is Regular , then $\exists V, V^* \in T$ such that $x \in V, V^* \supseteq w^{-1}\{1\}$ and $V \cap V^* = \emptyset$. But by the definition of lower semi continuous function $1_V, 1_{V^*} \in \omega(T)$ and $1_V(x) = 1, 1_{V^*} \geq 1_{w^{-1}\{1\}}, 1_V \cap 1_{V^*} = 1_{V \cap V^*} = 0$. Put $u = 1_V$ and $v = 1_{V^*}$, then , it is clear that $u(x) = 1, v \geq 1_{w^{-1}\{1\}}$ and $u \cap v \leq \alpha$.

Hence $(X, \omega(T))$ is α -FR(i).

It can easily be shown that $(2) \Rightarrow (3)$, $(3) \Rightarrow (5)$, $(2) \Rightarrow (4)$, $(4) \Rightarrow (5)$.

We therefore prove that $(5) \Rightarrow (1)$.

Let $(X, \omega(T))$ be α -FR(iv). We shall prove that (X, T) is Regular space. Let $x \in X$, V be a closed set in T , such that $x \notin V$. This implies that $V^c \in T$ and $x \in V^c$. But from the definition of $\omega(T)$, $1_{V^c} \in \omega(T)$, and $(1_{V^c})^c = 1_V$ closed in $\omega(T)$ and $1_V(x) = 0$. Since $(X, \omega(T))$ is α -FR(iv), for $\alpha \in I_1$, $\exists u, v \in \omega(T)$ such that $u(x) > \alpha$, $v \geq 1_{(1_V)^{-1}(1)} = 1_V$ and $u \cap v \leq \alpha$. Since $u, v \in \omega(T)$, then $u^{-1}(\alpha, 1], v^{-1}(\alpha, 1] \in T$ and $x \in u^{-1}(\alpha, 1]$. Since $v \geq 1_V$, then $v^{-1}(\alpha, 1] \supseteq (1_V)^{-1}(\alpha, 1] = V$, and $u \cap v \leq \alpha$ implies $(u \cap v)^{-1}(\alpha, 1] = u^{-1}(\alpha, 1] \cap v^{-1}(\alpha, 1] = \phi$. Now from the above it is clear that (X, T) is Regular space.

Thus it is seen that α -FR(p) is a good extension of its topological counterpart $(p = i, ii, iii, iv)$.

5.12. Theorem: - Let (X, t) and (Y, s) be two fuzzy topological spaces and

$f: X \longrightarrow Y$ be continuous, one-one, onto and open map then,

- (a) (X, t) is α -FR(i) $\Rightarrow (Y, s)$ is α -FR(i).
- (b) (X, t) is α -FR(ii) $\Rightarrow (Y, s)$ is α -FR(ii).
- (c) (X, t) is α -FR(iii) $\Rightarrow (Y, s)$ is α -FR(iii).
- (d) (X, t) is α -FR(iv) $\Rightarrow (Y, s)$ is α -FR(iv).

Proof :- Suppose (X, t) be α -FR(i). We shall prove that (Y, s) is α -FR(i). Let $w \in s^c$ and $p \in Y$ such that $w(p) < 1$, $f^{-1}(w) \in t^c$ as f is continuous and $x \in X$ such that $f(x) = p$ as f is one-one and onto. Hence $f^{-1}(w)(x) = w(f(x)) = w(p) < 1$. Since (X, t) is α -FR(i), for

$\alpha \in I_1$, then $\exists u, v \in t$ such that $u(x) = 1, v(y) = 1, y \in \{f^{-1}(w)\}^{-1}\{1\}$ and $u \cap v \leq \alpha$.

This implies that

$$f(u)(p) = \{ \text{Sup } u(x) : f(x) = p \} = 1.$$

and $f(v)(f(y)) = \{ \text{Sup } v(y) \} = 1$ as $f(f^{-1}(w)) \subseteq w \Rightarrow f(y) \in w^{-1}\{1\}$

and $f(u \cap v) \leq \alpha$ as $u \cap v \leq \alpha \Rightarrow f(u) \cap f(v) \leq \alpha$.

Now it is clear that $\exists f(u), f(v) \in s$ such that $f(u)(x) = 1, f(v)(f(y)) = 1, f(y) \in w^{-1}\{1\}$

and $f(u) \cap f(v) \leq \alpha$. Hence (Y, s) is α -FR(i).

Similarly (b), (c) and (d) can be proved.

5.13. Theorem:- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be a continuous, one-one, onto and closed map then,

(a) (Y, s) is α -FR(i) $\Rightarrow (X, t)$ is α -FR(i).

(b) (Y, s) is α -FR(ii) $\Rightarrow (X, t)$ is α -FR(ii).

(c) (Y, s) is α -FR(iii) $\Rightarrow (X, t)$ is α -FR(iii).

(d) (Y, s) is α -FR(iv) $\Rightarrow (X, t)$ is α -FR(iv).

Proof:- Suppose (Y, s) be α -FR(i). We shall prove that (X, t) is α -FR(i). $w \in t^\circ$ and $x \in X$ with $w(x) < 1$, then $f(w) \in s^\circ$ as f is closed and we find $p \in Y$ such that $f(x) = p$ as f is one-one. Now we have $f(w)(p) = \{ \text{Sup } w(x) : f(x) = p \} < 1$. Since (Y, s) is α -FR(i), for $\alpha \in I_1$, then $\exists u, v \in s$ such that $u(f(x)) = 1, v(y) = 1, y \in (f(w))^{-1}\{1\}$ and $u \cap v \leq \alpha$. This implies that $f^{-1}(u), f^{-1}(v) \in t$ as f is continuous and $u, v \in s$. Now $f^{-1}(u)(x) = u(f(x)) = u(p) = 1$ and $f^{-1}(v)(q) = v(f(q)) = v(y) = 1$ as $f(q) = y, y \in (f(w))^{-1}\{1\}$ ie $f(p) \in (f(w))^{-1}\{1\} \Rightarrow q \in w^{-1}\{1\}$ and $f^{-1}(u) \cap f^{-1}(v) = f^{-1}(u \cap v) \leq \alpha$ as $u \cap v \leq \alpha$. Now we observe that

$\exists f^{-1}(u), f^{-1}(v) \in \tau$ such that $f^{-1}(u)(x) = 1, f^{-1}(v)(q) = 1, q \in w^{-1}\{1\}$ and $f^{-1}(u) \cap f^{-1}(v) \leq \alpha$.

Hence (X, τ) is α -FR(i).

Similarly (b), (c) and (d) can be proved.

Chapter : - 6

Normal Fuzzy Topological Space

6. Introduction:-

In this chapter, we introduce and study Normal property in fuzzy topological spaces and obtain their several features.

6.1. Definition :- Let (X, τ) be a fuzzy topological space and $\alpha \in I_1$.

(a) (X, τ) is α - FN(i) space $\Leftrightarrow \forall w, w^* \in \tau^c$, with $w \cap w^* \leq \alpha$, $\exists u, u^* \in \tau$ such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* = 0$.

(b) (X, τ) is α - FN(ii) space $\Leftrightarrow \forall w, w^* \in \tau^c$ with $w \cap w^* = 0$, $\exists u, u^* \in \tau$ such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* \leq \alpha$.

(c) (X, τ) is a α - FN(iii) space $\Leftrightarrow \forall w, w^* \in \tau^c$ with $w \cap w^* \leq \alpha$, $\exists u, u^* \in \tau$ such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* \leq \alpha$.

6.2. Theorem :- The following implications are true:

(X, τ) is 0 - FN(i) $\Leftrightarrow (X, \tau)$ is 0 - FN(ii) $\Leftrightarrow (X, \tau)$ is 0 - FN(iii).

Proof:- Suppose (X, τ) be α - FN(i). We shall prove that (X, τ) is 0 - FN(ii). Let $w, w^* \in \tau^c$ with $w \cap w^* = 0$. This implies that $w \cap w^* \leq 0$. Since (X, τ) is 0 - FN(i), for $\alpha \in I_1$, $\exists u, u^* \in \tau$ such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* = 0$. It can be written as $u \cap u^* \leq 0$. Hence it is clear that (X, τ) is 0 - FN(ii).

Conversely, suppose that (X, t) is 0 - FN(ii) . We shall prove that (X, t) is 0- FN(i) .
 Let $w, w^* \in t^c$ with $w \cap w^* \leq 0$, ie $w \cap w^* = 0$. Since (X, t) is 0 -FN(ii), $\exists u, u^* \in t$
 such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$, and $u \cap u^* \leq 0$, ie $u \cap u^* = 0$.
 Hence it is clear that (X, t) is 0 - FN(i) .

Next, suppose that (X, t) is 0 - FN(ii) . We shall prove that (X, t) is 0 - FN(iii).
 Let $w, w^* \in t^c$ with $w \cap w^* \leq 0$, ie $w \cap w^* = 0$. Since (X, t) is 0 - FN(ii), $\exists u, u^* \in t$
 such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* \leq 0$. Hence it is clear
 that (X, t) is 0-FN(iii).

Conversely, suppose that (X, t) is 0 - FN(iii) . We shall prove that (X, t) is 0 - FN(ii).
 Let $w, w^* \in t^c$ with $w \cap w^* = 0$, ie $w \cap w^* \leq 0$. Since (X, t) is 0 - FN(iii), $\exists u, u^* \in t$
 such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$ and $u \cap u^* \leq 0$. Hence it is clear
 that (X, t) is 0-FN(ii).

6.3. Theorem :- If $0 \leq \alpha \leq \beta < 1$, then

(a) (X, t) is β - FN(i) $\Rightarrow (X, t)$ is α - FN(i) .

(b) (X, t) is α - FN(ii) $\Rightarrow (X, t)$ is β - FN(ii).

Proof :- First , suppose that (X, t) is β - FN(i) . We shall prove that (X, t) is α - FN(i) .
 Let $w, w^* \in t^c$ with $w \cap w^* \leq \alpha$. Since $0 \leq \alpha \leq \beta < 1$, then $w \cap w^* \leq \beta$. Since (X, t) is
 β - FN(i), for $\beta \in I_1, \exists u, u^* \in t$ such that $u(x) = 1, \forall x \in w^{-1}\{1\}, u^*(y) = 1, \forall y \in w^{*-1}\{1\}$
 and $u \cap u^* = 0$. Hence it is clear that (X, t) is a α - FN(i).

Next, suppose that (X, t) is α - FN(ii) . We shall prove that (X, t) is β - FN(ii) .
 Let $w, w^* \in t^c$ with $w \cap w^* = 0$. Since (X, t) is α -FN(ii), $\exists u, u^* \in t$ such that
 $u(x) = 1, x \in w^{-1}\{1\}, u^*(y) = 1, y \in w^{*-1}\{1\}$ and $u \cap u^* \leq \alpha$. Since $0 \leq \alpha \leq \beta < 1$, then
 $u \cap u^* \leq \beta$. Hence it is clear that (X, t) is β - FN(ii) .

6.4. Theorem :- Let (X, t) be a fuzzy topological space, and

$I_\alpha(t) = \{ u^{-1}(\alpha, 1] : u \in t \}$, then

(X, t) is 0-FN(iii) space $\Rightarrow (X, I_0(t))$ is Normal space.

Proof :- Suppose that (X, t) be a α -FN(iii) space. We shall prove that $(X, I_0(t))$ is Normal space. Let V, V^* be closed set in $I_0(t)$, and $V \cap V^* = \phi$. Then $V^c, V^{*c} \in I_0(t)$ and $(V \cap V^*)^c = V^c \cup V^{*c} = X$. Since $V^c, V^{*c} \in I_0(t)$, then, $\exists u, u^* \in t$ such that $V^c = u^{-1}(0, 1]$ and $V^{*c} = u^{*-1}(0, 1]$ and $u^{-1}(0, 1] \cup u^{*-1}(0, 1] = V^c \cup V^{*c} = X$. Hence $(u \cup u^*)^{-1}(0, 1] = X$. Now we find $u^c, u^{*c} \in t^c$ such that $((u \cup u^*)^{-1}(0, 1])^c = \phi$. This implies that $(u \cup u^*)^c = u^c \cap u^{*c} = 0$. Since (X, t) is 0-FN(iii), $\exists v, v^* \in t$ such that $v \geq 1_{(u^c)^{-1}\{0\}}$, $v^* \geq 1_{(u^{*c})^{-1}\{0\}}$, $v \cap v^* = 0$. But from the definition of $I_0(t)$, $v^{-1}(0, 1], v^{*-1}(0, 1] \in I_0(t)$, and we get $v^{-1}(0, 1] \supseteq 1_{(u^c)^{-1}\{0\}}(0, 1], v^{*-1}(0, 1] \supseteq 1_{(u^{*c})^{-1}\{0\}}(0, 1], (v \cap v^*)^{-1}(0, 1] = \phi$. Put $W = v^{-1}(0, 1], W^* = v^{*-1}(0, 1]$ in $I_0(t)$. Then finally we find, $W \supseteq V, W^* \supseteq V^*$ and $W \cap W^* = \phi$. Hence it is clear that $(X, I_0(t))$ is Normal space.

6.5. Theorem :- Let (X, T) be a topological space consider the following statements:

- (a) (X, T) be a Normal space
- (b) $(X, \omega(T))$ be α -FN(i) space.
- (c) $(X, \omega(T))$ be α -FN(ii) space.
- (d) $(X, \omega(T))$ be α -FN(iii) space.

Then (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (a).

Proof :- First suppose that (X, T) be a Normal space. We shall prove that $(X, \omega(T))$ be α -FN(i) space. Let w, w^* be closed in $\omega(T)$ and $w \cap w^* \leq \alpha$. Then it is clear that $w^c, w^{*c} \in \omega(T)$ and $(w \cap w^*)^c \geq 1 - \alpha > 0$. But from the definition of $\omega(T)$, $(w^c)^{-1}(0, 1],$

$(w^*)^{-1}(0, 1] \in T$. Now we see that $((w \cap w^*)^c)^{-1}(0, 1] = X$, and we also see that $((w^c)^{-1}(0, 1])^c = w^{-1}\{1\}$ and $((w^*)^c)^{-1}(0, 1])^c = (w^*)^{-1}\{1\}$ be closed in T . Now $((w \cap w^*)^c)^{-1}(0, 1])^c = (w \cap w^*)^{-1}\{1\} = w^{-1}\{1\} \cap (w^*)^{-1}\{1\} = \phi$. Since (X, t) is fuzzy Normal, then, $\exists V, V^* \in T$ such that $V \supseteq w^{-1}\{1\}, V^* \supseteq (w^*)^{-1}\{1\}$ and $V \cap V^* = \phi$. But from the definition of $w(T), 1_V, 1_{V^*} \in w(T), 1_V \geq 1_{w^{-1}\{1\}}, 1_{V^*} \geq 1_{(w^*)^{-1}\{1\}}$ and $1_{V \cap V^*} = 0$.

Hence $(X, w(T))$ is α -FN(i) space.

It can easily be shown that $(b) \Rightarrow (c) \Rightarrow (d)$.

We, therefore prove that $(d) \Rightarrow (a)$.

Suppose, that $(X, w(T))$ is α -FN(iii). We shall prove that (X, T) is Normal space.

Let $V, V^* \in T^c$ and $V \cap V^* = \phi$. Then it is clear that $1_V, 1_{V^*}$ be closed in $w(T)$ and $1_{V \cap V^*} = 0$. Since $(X, w(T))$ is α -FN(iii), then $\exists u, u^* \in w(T)$ such that $u \geq 1_V, u^* \geq 1_{V^*}$ and $u \cap u^* \leq \alpha$. But from the definition of $w(T), u^{-1}(\alpha, 1], (u^*)^{-1}(\alpha, 1] \in T$ and $u^{-1}(\alpha, 1] \supseteq (1_V)^{-1}(\alpha, 1] = V, (u^*)^{-1}(\alpha, 1] \supseteq (1_{V^*})^{-1}(\alpha, 1] = V^*$ and $u^{-1}(\alpha, 1] \cap (u^*)^{-1}(\alpha, 1] = (u \cap u^*)^{-1}(\alpha, 1] = \phi$. Hence it is clear that (X, T) is Normal space.

Thus it is seen that α -FN(p) is a good extension of its topological counter part.
 . (p = i, ii, iii)

6.6. Theorem :- Let (X, t) and (Y, s) be two fuzzy topological spaces and $f: X \longrightarrow Y$ be a continuous, one-one, onto and open map then,

- (a) (X, t) is α -FN(i) $\Rightarrow (Y, s)$ is α -FN(i).
- (b) (X, t) is α -FN(ii) $\Rightarrow (Y, s)$ is α -FN(ii).
- (c) (X, t) is α -FN(iii) $\Rightarrow (Y, s)$ is α -FN(iii).

Proof :- Suppose (X, t) be α -FN(i). We shall prove that (Y, s) is α -FN(i). Let $w, w^* \in s^c$ with $w \cap w^* \leq \alpha$ then $f^{-1}(w), f^{-1}(w^*) \in t^c$ as f is continuous. Now $f^{-1}(w \cap w^*) \leq \alpha$ as $w \cap w^* \leq \alpha \Rightarrow f^{-1}(w) \cap f^{-1}(w^*) \leq \alpha$. Since (X, t) is α -FN(i), for $\alpha \in I_1$, then $\exists u, u^* \in t$ such that $u(x) = 1, x \in (f^{-1}(w))^{-1}\{1\}, u^*(y) = 1, y \in (f^{-1}(w^*))^{-1}\{1\}$ and $u \cap u^* = 0$. This implies that $f(u), f(u^*) \in s$ as f is open.

Now $f(u)(p) = \{ \text{Sup } u(x) \ ; \ f(x) = p \}, f^{-1}(p) \in (f^{-1}(w))^{-1}\{1\}$

ie $f(u)(p) = 1, p \in w^{-1}\{1\}$

and $f(u^*)(q) = \{ \text{Sup } u^*(y) \ ; \ f(y) = q \}, f^{-1}(q) \in (f^{-1}(w^*))^{-1}\{1\}$

ie $f(u^*)(q) = 1, q \in w^{*-1}\{1\}$ and $f(u) \cap f(u^*) = f(u \cap u^*) = 0$ as $u \cap u^* = 0$.

Now it is clear that $\exists f(u), f(u^*) \in s$ such that $f(u)(p) = 1, p \in w^{-1}\{1\}, f(u^*)(q) = 1, q \in w^{*-1}\{1\}$ and $f(u) \cap f(u^*) = 0$. Hence (Y, s) is α -FN(i).

Similarly (b) and (c) can be proved.

6.7. Theorem :- Let (X, t) and (Y, s) be two fuzzy topological spaces and

$f: X \longrightarrow Y$ be continuous, one-one, onto and closed map then,

(a) (Y, s) is α -FN(i) $\Rightarrow (X, t)$ is α -FN(i).

(b) (Y, s) is α -FN(ii) $\Rightarrow (X, t)$ is α -FN(ii).

(c) (Y, s) is α -FN(iii) $\Rightarrow (X, t)$ is α -FN(iii).

Proof:- Suppose (Y, s) be α -FN(i). We shall prove that (X, t) is α -FN(i). Let $w, w^* \in t^c$ with $w \cap w^* \leq \alpha$, then $f(w), f(w^*) \in s^c$ as f is closed and $f(w) \cap f(w^*) = f(w \cap w^*) \leq \alpha$, as $w \cap w^* \leq \alpha$. Since (Y, s) is α -FN(i), for $\alpha \in I_1$, then $\exists u, u^* \in s$ such that $u(x) = 1, x \in (f(w))^{-1}\{1\}, u^*(y) = 1, y \in (f(w^*))^{-1}\{1\}$ and $u \cap u^* = 0$. This implies that $f^{-1}(u), f^{-1}(u^*) \in t$ as f is continuous and $u, u^* \in s$.

Now $f^{-1}(u)(p) = u(f(p)) = u(x) = 1$ as $f(p) = x \in (f(w))^{-1}\{1\}$

ie $f^{-1}(u)(p) = 1, p \in w^{-1}\{1\}$

$f^{-1}(u^*)(q) = u^*(f(q)) = u^*(y) = 1$ as $f(q) = y \in (f(w^*))^{-1}\{1\}$

ie $f^{-1}(u^*)(q) = 1, q \in w^{*-1}\{1\}$

and $f^{-1}(u) \cap f^{-1}(u^*) = f^{-1}(u \cap u^*) = 0$.

Now it is clear that $\exists f^{-1}(u), f^{-1}(u^*) \in \tau$ such that $f^{-1}(u)(p) = 1, p \in w^{-1}\{1\}, f^{-1}(u^*)(q) = 1, q \in w^{*-1}\{1\}$ and $f^{-1}(u) \cap f^{-1}(u^*) = 0$. Hence (X, τ) is α -FN(i).

Similarly (b) and (c) can be proved.

References

1. Ajmal, N.; Tyagi , B. K. : A Characterization of regular fuzzy space; Mat. Vesnik 41(1989) no. 2, 65-70.
2. Ali, D. M. : A note on fuzzy regularity concepts; Fuzzy sets and systems. 35(1990), 101-104.
3. Ali , D. M. ; Wuyts, P. ; Srivastava, A. K. : On the R_0 – property in fuzzy topology; Fuzzy sets and systems. 38(1990) no. 1, 97-113.
4. Ali , D. M. : A note on some FT_2 concepts; Fuzzy sets and systems. 42(1991) no. 3, 381-386.
5. Ali , D. M. : Some other types of fuzzy connectedness; Fuzzy sets and systems. 46(1992), 55-61.
6. Ali , D. M. : On the R_1 - property in fuzzy topology; Fuzzy sets and systems. 50(1992), 97-101.
7. Allam , A. A.; Abt. E1, K. M. ; Nehad N. Morsi: On Fuzzy neighbourhood spaces: Fuzzy sets and systems. 41 (2) (1991) 201-212.
8. Arya, S. P. ; Nour, T. M. : Separation axioms for bi -topological spaces; Jour. Pure Appl. Math. 19(1) (1988) 42-50.
9. Azad, K. K. : On Fuzzy semi- continuity, Fuzzy almost continuity and Fuzzy weakly continuity; J. Math. Anal . Appl. 82(1) (1981) 14-32.
10. Bennet , G.; Kalton, N. J. : FK-spaces containing co; Duke Math Jour. 39(1972), 561-582.
11. Bhaumik, R. N. ; Mukherjee, A. : Some more results on completely induced fuzzy topological spaces, Fuzzy sets and systems. 50 (1992) 113-117.
12. Buck, R. C. : Operator algebras and dual spaces. Proc. Amer. Math. Soc. 3(1952), 681-687.

13. Carmer , J. H. : L – topologische Raume (un published) Universitat Bremen
(1986) .
14. Chakravarty, K. K.; Ahsanullah, T. M. G. : Fuzzy topology on fuzzy sets and
tolerance topology; Fuzzy sets and systems. 45 (1992) 103-108.
15. Chang, C. L. : Fuzzy topological spaces; J. Math. Anal Appl. 24(1968), 182-192.
16. Chattopadhyya, K. C. ;Hazra, R. N.; Samanta, S. K. : Gradation of openness fuzzy
topology; Fuzzy sets and systems. 49 (1992) 1237-242.
17. Cooper, J. B. : The strict topology and space with mixed topologies; Proc. Amer.
Math. Soc. 30 (1971),583-592.
18. Cooper, J. B. : Saks space and applications to Functional Analysis. North Holland
Publishers 1978.
19. Cutler, D. R. ; Reilly, I. L : A comparison of some Hausdorff notions in fuzzy
topological spaces; Comput. Math. Appl. 19(1990) no. 11, 97-104.
20. Das, N. R. ; Das , P. : Mixed topological groups. Indian J. Pure appl. Math. 22(4)
(1991), 323-329.
21. Das, N. R. ; Baishya, P. C. : Mixed fuzzy topological spaces. The Journal of Fuzzy
Math. Vol. 3 No. 4 December 1995.
22. Das, N. R. ; Baishya, P. C. : On open maps, closed maps and fuzzy continuous
maps in a fuzzy bi-topological spaces. (Communicated)
23. Dube, K. K. ; Misra, D. N. : Some localized separation axioms and their
applications; M. R. 59(3) (1979)
24. Dubois, D. ; Prade, H. : Fuzzy sets and systems. Theory and Applications;
Academic Press New York 1980.
25. Erceg, M. A. : Functions Equivalence relations, Quotient spaces and subjects in
Fuzzy set theory; Fuzzy sets and systems. 3(1) (1980) , 75-92.

26. Eroglue, M. S. : Topological representation for fuzzy topological spaces; Fuzzy sets and systems. 42(1991), 335-362.
27. Evert, J. : On normal fuzzy topological spaces; Mathematica (cluj) 31(54) (1989) no. 1, 39-45.
28. Fath Alla, M. A. ; Abd El . Hakeim, K. M. : Mapping with fuzzy closed graphs and strongly fuzzy closed graphs; J. Inst. Math. Comput. Sci. Math. Ser 4(1991) no. 1, 61-67.
29. Fora, Ali Ahmed : Separation axioms, subspace and product spaces in fuzzy topology; Arab Culf J. Sci. Res. 8(1990) no.3, 1-16.
30. Foster, D. H. : Fuzzy topological groups; J. Math. Anal. Appl. 67(1979), 549-567.
31. Ganguly, S ; Saha, S. : On separation axioms and separations of connected sets in fuzzy topological spaces.; Bull.Cal. Math. Soc. 79(1987), 215-225.
32. Ghanim, M. H. ; Kerre, E. E. ;Mashhour, A. S. : Separation axioms, subspace and sums in fuzzy topology; J. Math. Anal. Appl. 102(1984), 189-202.
33. Ghanim, M. H. ; Morsi, N. N. : α - axims in fuzzy topology; Simon Stevin 63(1989) no. 3-4, 193-208.
34. Ghanim, M. H. : Pseudo- closure operators in fuzzy topological spaces ; Fuzzy sets and systems. 39(1991) no. 3, 339-346.
35. Goguen, J. A. : L-fuzzy sets; J. Math. Anal. Appl. 18(1967), 145-174.
36. Goguen, J. A. : The fuzzy Tychonoff Theorem; J. Math. Anal. Appl. 43(1973), 734-742.
37. Hazra, R. N. ; Samanta, S. K.; Chattopadhyya : Fuzzy topology redefined ; Fuzzy sets and systems. 45(1992) , 79-82.
38. Hutton, B ; Reilly, I. L. : Separstion axioms in fuzzy topological spaces; Dept. Math. Univ. of Auckland, Report No. 55, March 1974.

39. Hutton, B. : Normality in fuzzy topological spaces; J. Math. Anal. Appl. (1975), 74-79.
40. Hutton, B.: Uniformities on fuzzy topological spaces; J. Math. Anal. Appl. 58(1977), 559-571.
41. Hutton, B.: Products of fuzzy topological spaces Topology; Appl. 11(1980), 59-67.
42. Katsaras, A. K. : Fuzzy neighborhood structures and fuzzy quasi uniformities; Fuzzy sets and systems. 29(1989) no. 2, 187-199.
43. Kaufmann, A.: Introduction to the theory of fuzzy subsets. Volume I; Academic Press , I. N. C. (1975).
44. Kaufmann, A.; Gupta, M. M.: Introduction to fuzzy Arithmetic Theory and Application. Van Nostrand Reinhold Company. New York.
45. Kelley, J. L. : General Topology ; Van Nostrand, Princeton, NJ, 1955.
46. Kerre, E. E. ; Ottey, P. L. : On α -generated fuzzy topologies; Fasc. Math. No. 19 (1990), 127-134.
47. Lipschutz , S . : General , Topology ; Copyright © 1965 , by the Schaum publishing Company .
48. Lowen, R.: Topologies floues; C. R. Acad Sc. Parish 278 series A(1974), 925-928.
49. Lowen, R.: Initial and final topologies and fuzzy Tychonoff Theorem; J. Math. Anal. Appl. 58(1977), 11-21.
50. Lowen, R.: Fuzzy topological spaces and fuzzy compactness; J. Math. Anal. Appl. 56(1976), 621-633.
51. Macho, Stadler, M.; De Prada Vicente M. A. : Strong separation and strong countability in fuzzy topological spaces; Fuzzy sets and systems. 43(1991) no. 1, 95-116.
52. Malghan, S. R. ; Benchalli, S. S. : On fuzzy topological spaces; Glasnic Mathematicki, 16(36) (1981), 313-325.

53. Malghan, S. R. ; Benchalli, S. S. : On open Maps , closed Maps and Local compactness in fuzzy topological spaces; J. Math. Anal. Appl. 99(2) (1984), 338-349.
54. Mashhour, A. S. ; Morsi, Nehad, N; El. Tantawy, O. A. : Fuzzy neighborhood syntopogenous structure; Fuzzy sets and systems. 43(1991) no. 2 , 219-234.
55. Mashhour, A. S. ; Morsi, N, N. : On regularity axioms in fuzzy neighborhood spaces; Fuzzy sets and systems. 44(1991) , 265-271.
56. Ming, Pu. Pao ; Ming, Liu Ying : Fuzzy topology I. Neighborhood Structure of a fuzzy point and Moore - Smith convergence; J. Math. Anal. Appl. 76 (1980), 571-599.
57. Ming, Pu. Pao ; Ming, Liu Ying : Fuzzy topology II. Product and Quotient Spaces; J. Math. Anal. Appl. 77 (1980), 20-37.
58. Ming Sheng Ying : A new approach for fuzzy topology I; Fuzzy sets and systems. 39(1991) , 303-321.
59. Ming Sheng Ying : A new approach for fuzzy topology II; Fuzzy sets and systems. 49(1992) , 221-232.
60. Morsi, Nehad N.: The Urysohn Lemma for fuzzy neighbourhood spaces; Fuzzy sets and systems. 39(1991) no. 3, 347-360.
61. Morsi, Nehad N.: Dual fuzzy neighbourhood spaces I; Fuzzy sets and systems. 44(1991) , 245-263.
62. Mukherjee, M. N. ;Sinha, S. P. : On some near fuzzy countinuous functions between fuzzy topological spaces ; Fuzzy sets and systems. 34 (1990) no. 2, 245-254.
63. Mukherjee, M. N. ; Ghosh, B. : Fuzzy semi- regularzation topologies and Fuzzy submaximal spaces; Fuzzy sets and systems. 49(1991) , 283-294.

64. Nanda, S. ; S. Dang , A. Behera : The fuzzy topological complementation Theorem;
Jour. Fuzzy Math, 1(2) (1993), 303-310.
65. Patronis, T.; Stravrinos,P. : Fuzzy equivalence and the resulting topology; Fuzzy sets
and systems. 46(1992) , 237-243.
66. Petricevic , zlata : R_0 and R_1 axioms in fuzzy topology; Mat. Vesnik 41(1989),21-
28.
67. Petricevic , zlata : Some separation axioms in fuzzy spaces; zb, Rad. No. 4(1990) 25
– 33 .
68. Ramadan, A. A.; Abd Alla , M. A. : On smooth pre-proximity spaces; Fuzzy sets and
systems. 48(1992) , no 1 , 117 – 121 .
69. Ramadan, A. A.: Smooth topological spaces; Fuzzy sets and systems. 48(1992) ,
371-375.
70. Rodabaugh , S. E. ; The Hausdorff Separation axiom for fuzzy topological spaces.
Top. Appl (1980) , 319 – 334 .
71. Rudin , W.; Real and complex analysis . Copyright © 1966 , 1974 , by McGraw –
hill Inc .
72. Sentilles , F. Dannis : Bounded continuous functions on a completely regular space;
Trans . Amer. Math. Soc. Vol. 168(1972).
73. Shostak, A. P. : Two decades of fuzzy topology : the main ideas , concepts and
results. (Russian). Uspekhi. Mat. Nauk 44(1989) 99-147.
74. Singal, M. K. ; Prakash, N. : Fuzzy pre-open sets and fuzzy pre separation axioms;
Fuzzy sets and systems. 44(1991) , 273-281.
75. Singal, M. K. ; Rajvanshi, N. : Regularly open sets in fuzzy topological spaces;
Fuzzy sets and systems. 50(1992) , 343-353.
76. Sinha, S. P. : Fuzzy normality and some of its weaker forms; Bull. Korean. Math.
Soc. 28(1991) no. 1, 89-97.

77. Sinha, S. P. : Separation axioms in fuzzy topological spaces; Fuzzy sets and systems. 45(1992) , 261-270.
78. Sostak, A. P. : On the neighborhood structure of fuzzy topological spaces; Zb, Rad. No. 4(1990) , 7-14.
79. Sostak , Alexander; Dzhajanbajev : On completeness and pre-compactness spectra of fuzzy sets in fuzzy uniform spaces; Comment. Math. Univ. Car. 31 (1990) no. 1,149-157.
80. Srivastava , R.; Lal, S. N. ; Srivastava, A. K : Fuzzy T_1 - topological spaces; J. Math . Anal. Appl.102 (1984), 442-4448.
81. Srivastava , A. K.; Ali , D. M. : A note on K. K. Azad's fuzzy hausdorffness concepts; Fuzzy sets and Systems. 42 (1991) , 363-367.
82. Stadler, M. M. ; De prada Vicente M. A. : Strong separation and strong countability in fuzzy topological spaces ; Fuzzy sets and System. 43 (1991) no. 1, 95-116.
83. Warner , M. W. ; Mclean , R. G. : On compact Hausdorff L-fuzzy spaces ; Fuzzy Sets and Systems. 56 (1993) , 103-110.
84. Warren , R. H. : Neighbourhoods, bases and continuity in fuzzy topological spaces ; Rocky Mountain J. Math. 8 (1978). 459-470.
85. Warren , R. H. : Fuzzy topologies characterized by neighborhood systems ; Rocky Mountain J. Math. 9 (4) (1997), 761-764.
86. Weiss, M. D. : Fixed points , separation and induced topologies for fuzzy sets ; J. Math . Anal . Appl . 50 (1975), 142-150.
87. Wong , C. K. : Covering properties of fuzzy topological spaces ; J. Math . Anal. Appl. 43 (1973), 697-704.
88. Wong , C. K. : Fuzzy topology : Product and Quotient Theorem . J. Math . Anal . Appl. 45 (1974) 512-521.

89. Wong , C. K. : Fuzzy points and local properties of Fuzzy topology ; J. Math. Anal . Appl. 46 (1974), 316-328.
90. Wuyts, P.; Lowen , R.: On separation axioms in fuzzy topological spaces ; Fuzzy neighborhood spaces and Fuzzy Uniform spaces ; J . Math. Anal . Appl . 93 (1973) , 27 – 41 .
91. Xu, Luo Shan : The strong Hausdorff Separation property in L- fuzzy topological spaces ; Mohu. Xitong . Yu . Shuxue 5 (1991) no. 2, 25-29.
92. Ying , Ming Sheng : A new approach for fuzzy topology I ; Fuzzy Sets and Systems. 39 (1991) no. 3, 303-321.
93. Zhang , Ghong Quan : Fuzzy continuous function and its properties ; Fuzzy Sets and Systems. 43 (1991), 159-171.
94. Zadeh, L. A. : Fuzzy sets. Information and control 8 (1965) . 338-353.

Rajshahi University Library
Document Section
Document No. D-2436
Date... 2/10/05